

# Systematic Review of the Technique of Colorectal Anastomosis

Juliette C. Slieker, MD; Freek Daams, MD; Irene M. Mulder, MD;  
Johannes Jeekel, MD, PhD; Johan F. Lange, MD, PhD

**M**any different techniques of colorectal anastomosis have been described in search of the technique with the lowest incidence of anastomotic leak. A systematic review of leak rates of techniques of hand-sewn colorectal anastomosis was conducted to provide a guideline for surgical residents and promote standardization of its technique. Clinical and experimental articles on colorectal anastomotic techniques and anastomotic healing published in the past 4 decades were searched. We included evidence on suture material, suture format, single- vs double-layer sutures, interrupted vs continuous sutures, hand-sewn vs stapled and compression colorectal anastomosis, and anastomotic configuration. In total, 3 meta-analyses, 26 randomized controlled trials, 11 nonrandomized comparative studies, 20 cohort studies, and 57 experimental studies were found. Results show that, for many aspects of the hand-sewn colorectal anastomosis technique, evidence is lacking. A single-layer continuous technique using inverting sutures with slowly absorbable monofilament material seems preferable. However, in contrast to stapled and compression colorectal anastomoses, the technique for hand-sewn colorectal anastomoses is nonstandardized with regard to intersuture distance, suture distance to the anastomotic edge, and tension on the suture. We believe detailed documentation of the anastomotic technique of all colorectal operations is needed to determine the role of the hand-sewn colorectal anastomosis.

*JAMA Surg.* 2013;148(2):190-201

Construction of a colorectal anastomosis is a hallmark of surgical training. However, although surgical residents can refer to key publications with evidence-based conclusions for many topics, mere imitation of an experienced surgeon traditionally is considered the basic source for the technique of hand-sewn colorectal anastomosis. The large variety of anastomotic techniques is one of the main difficulties in the interpretation of the literature. Anastomotic leak (AL) following colorectal resection is a major problem of surgical care, with an incidence between 3% and 19%.<sup>1-4</sup> Although accurate prediction of risk is impossible, certain factors are known to contribute to AL, in-

cluding surgeon-related factors (eg, increased incidence of AL in a colorectal anastomosis constructed after hours<sup>5</sup> and the positive role of specialization in reducing the complications of colorectal surgery<sup>6</sup>) and patient-related risk factors (eg, the inverse relationship between the height of the colorectal anastomosis from the anal verge and the leak rate<sup>7-12</sup>). Decades of research have resulted in many studies investigating different techniques for constructing colorectal anastomosis in search of the safest method. Appreciating the conclusions from this extensive research is essential for the quality of colorectal surgery and for the resident being trained in colorectal surgery. Our aims were to perform a systematic review of all aspects of the technique of hand-sewn colorectal anastomosis and compare hand-sewn with mechanical colorectal anastomosis to pro-

**Author Affiliations:** Departments of Surgery (Drs Slieker, Daams, Mulder, and Lange) and Neuroscience (Dr Jeekel), Erasmus Medical Center, Rotterdam, the Netherlands.

vide a guideline for residents and promote standardization of the technique.

## METHODS

### SEARCH STRATEGY

A literature search was conducted using MEDLINE, EMBASE, and Cochrane databases for studies published between January 1, 1970, and February 1, 2011, using the key words presented in the eFigure (<http://www.jamasurg.com>). The search was restricted to articles published in English, Dutch, and French. References in the selected publications were searched for additional studies.

### STUDY SELECTION

Clinical as well as experimental studies were selected to address several aspects of the technique of hand-sewn colorectal anastomosis. These included:

1. suture material,
2. suture format (size of suture bites, in-between distance of bites, suture tension, configuration of the bite, and inverting vs everting sutures),
3. single- vs double-layer colorectal anastomosis,
4. interrupted vs continuous sutures,
5. hand-sewn vs stapled colorectal anastomosis,
6. hand-sewn vs compression colorectal anastomosis, and
7. configuration of colorectal anastomosis (end-to-end [ETE], end-to-side, side-to-end, side-to-side, length of the side-limb, and length of the enterotomy).

### INCLUSION CRITERIA FOR CLINICAL STUDIES

Only clinical studies comparing 2 or more colorectal anastomotic techniques with regard to clinical AL were considered relevant. When only 1 comparative study was available on a particular subject, clinical cohort studies were added to the selection. Results were analyzed only if the study groups and results were clearly described with proper statistical analysis.


### INCLUSION CRITERIA FOR EXPERIMENTAL STUDIES

Experimental studies were selected when comparing 2 or more colorectal anastomotic techniques using objective measurements for anastomotic healing: AL, anastomotic bursting pressure (ABP), anastomotic breaking strength, histologic results, or collagen concentration.

When 2 studies were reported by the same institution, either the better quality study or the most recent publication was included. As with clinical studies, results were analyzed only if the study groups and results were clearly described with proper statistical analysis. However, the lack of statistical analysis of histologic findings in experimental studies was accepted.

### EXCLUSION CRITERIA

Because the healing of small-bowel anastomoses is different and the incidence of AL is lower compared with large-bowel anastomoses, studies including both procedures without differentiating the results and statistical analysis were excluded. Ileocolic anastomoses after right hemicolectomy or ileocecal resection represent healing of the colon and were therefore



**Figure.** The PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analyses) flowchart: selection of relevant studies.

included. Studies reporting radiologic AL without distinction of clinical AL were excluded, as were studies reporting only on emergency operations, children, and colo-anal anastomosis or pouches. Results of experimental studies measured directly after the construction of colorectal anastomosis were not taken into account because these do not reflect anastomotic healing.

### DATA EXTRACTION FOR CLINICAL STUDIES

Two physicians (J.C.S. and F.D.) entered data in a database following standard protocols. Seven factors were considered for clinical studies. These included:

- first author and year of publication,
- level of evidence (following the Centre of Evidence Based Medicine, University of Oxford),
- study design,
- number of patients,
- location of anastomosis in the gastrointestinal tract,
- definition of outcome by the authors (AL, clinical AL, and radiologic AL), and
- results and statistical analysis.

### DATA EXTRACTION FOR EXPERIMENTAL STUDIES

Six factors were considered for experimental studies. These included:

- first author and year of publication,
- study design,
- number of animals per group,
- species,
- outcome factors for anastomotic healing (AL, ABP, breaking strength, histologic results, or collagen concentration), and
- results.

## RESULTS

The literature search identified 6168 articles; 1443 articles remained after duplicates were removed. The PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analyses) flowchart presented in the **Figure** shows the selection of studies: 117 were included in the systematic review. Included studies

**Table 1. Included Studies on Suture Material**

Source	Level <sup>a</sup>	No. of Cases	Outcome	Results
<b>Clinical</b>				
Gillatt et al, <sup>24</sup> 1987	1b (RCT)	57 Patients, colorectal	Clinical AL	AL: PDS, 0 (n = 30); silk, 3.7% (n = 27); not significant
Clark et al, <sup>23</sup> 1977	1b (RCT)	194 Patients, colon	Clinical and radiologic AL	AL: catgut, 24% (n = 99); polyglycolic acid, 8.4% (n = 95); significant
<b>Experimental</b>				
Andersen et al, <sup>20</sup> 1989	...	196 Rats	Histologic results	Polyglactin 910 vs PDS: equal histologic results at days 7 and 56
Foresman et al, <sup>17</sup> 1989	...	160 Rats	ABP	Polyglyconate vs PDS: difference in ABP at days 0, 7, 14, 21, 42 not significant
Lord et al, <sup>18</sup> 1978	...	30 Rats	Histologic results of the submucosa	Rough surface vs smooth surface: rough surface (catgut, braided silk, polyglycolic acid), most damage to submucosa; polyethylene terephthalate coated with polytetrafluoroethylene better, polypropylene suture best histologic results
Orringer et al, <sup>15</sup> 1977	...	84 Dogs	Histologic results	Silk vs polypropylene vs wire: silk, most inflammation; polypropylene, less inflammation; wire, least inflammation at days 4, 7, 10, 14
Munday and McGinn, <sup>19</sup> 1976	...	44 Rats	ABP, histologic results	Catgut vs polyglycolic acid: equal ABP and histologic results at day 7
Deveney and Way, <sup>16</sup> 1977	...	60 Dogs	Breaking strength, histologic results	Catgut vs chromic catgut vs polyglycolic acid vs polyglactin 910: equal strength and equal histologic results at days 4, 7, 14
Hastings et al, <sup>13</sup> 1975	...	127 Dogs	Breaking strength, histologic results, collagen	Absorbable vs nonabsorbable sutures: absorbable, less breaking strength at days 14 and 28, equal collagen at day 120, worse histologic results at day 120
Letwin, <sup>14</sup> 1975	...	28 Dogs	Histologic, results, ABP	Double-layer anastomoses "chromic catgut-silk" vs polyglycolic acid-polyglycolic acid: catgut-silk, significantly more suture reaction and significantly lower ABP at day 7
Pasternak et al, <sup>22</sup> 2008	...	85 Rats	Breaking strength	Doxycycline (MMP-i)-coated sutures vs control carrier-coated sutures: higher breaking strength in doxycycline sutures at days 0-3
Pascual et al, <sup>21</sup> 2008	...	40 Rats	AL, ABP	Polyglactin 910 with mesenchymal cells vs polyglactin 910: no difference in AL, no difference in ABP; fewer adhesions in polyglactin 910 with mesenchymal cells at days 4, 7, 14, 21

Abbreviations: ABP, anastomotic bursting pressure; AL, anastomotic leak; MMP-i, matrix metalloproteinase inhibitor; PDS, polydioxanone sutures; RCT, randomized controlled trial; ellipses indicate not applicable.

<sup>a</sup>Highest level of evidence, 1b.

and their characteristics are listed in **Tables 1, 2, 3, 4, 5, 6,** and **7**, together with all results of outcome measures. The results per research question are summarized herein.

## SUTURE MATERIAL

Decades ago, several materials, such as silk, linen, catgut, polyglactin 910, and nylon, were commonly used for colorectal anastomosis. Today most gastrointestinal anastomoses, including colorectal anastomosis, are constructed with polydioxanone sutures. Ten experimental studies<sup>13-22</sup> were included. Results show that absorbable sutures compared with nonabsorbable or slowly absorbable sutures cause more tissue reaction<sup>13-15</sup>; one of these studies<sup>13</sup> showed that absorbable sutures dissolve too rapidly, influencing anastomotic strength. Multifilament compared with monofilament sutures cause more tissue damage and easier adherence of material within the interstices of multifilament sutures,<sup>16-19</sup> providing a basis for infection.<sup>121</sup> Surprisingly, experimental studies on the healing of colorectal anastomosis constructed with polydioxanone sutures are scarce; only 2 studies<sup>17,20</sup> were included, finding equal ABP and histologic characteristics between polydioxanone and polyglycolic acid. Noncomparative experimental studies<sup>121-124</sup> that did not meet the inclusion criteria for the present systematic review have shown that polydioxanone sutures possess all aspects considered important: monofilament, little histologic reaction, slowly absorbable with long preservation of strength, and low adherence of bacteria to the material.

New possibilities for the use of sutures coated with mesenchymal stem cells and doxycycline were explored

in 2 experimental studies with promising, but not yet convincing, results.<sup>21,22</sup> Two included randomized controlled trials (RCTs)<sup>23,24</sup> on suture material failed to achieve a unanimous conclusion because of the small number of patients included and the different suture materials tested that are rarely used today.

In conclusion, on the basis of experimental studies, nonabsorbable or slowly absorbable monofilament sutures seem to be the first choice for colorectal anastomosis. However, there is no level 1 evidence to confirm this hypothesis.

## SUTURE FORMAT

### Size of Suture Bites

Since Lembert<sup>125</sup> described the construction of intestinal anastomoses in dogs using suture bites with 5-mm distance to the cut edge nearly 2 centuries ago, this aspect seems to have become less clear in surgical literature. One experimental study<sup>25</sup> was found for this systematic review that investigated the difference in anastomotic strength in rats with sutures placed between 3 mm and 1.5 mm from the cut edges. Results showed lower breaking strength for small bites, measured at day 2. One RCT by Greenall et al<sup>26</sup> reporting on the distance of the suture to the wound edge matched the inclusion criteria. They randomly allocated patients to have bowel sutures placed either 5 or 10 mm from the cut edges, with no significant differences in AL. Because it is not possible to extrapolate the distances used in a rat model to the clinical situation, we can only conclude from one level 1b RCT that distances of 5 and 10 mm from the cut edge will probably give adequate results.

**Table 2. Included Studies on Suture Format**

Source	Level	No. of Cases	Outcome	Results
<b>Size of suture bites<sup>a</sup></b>				
Clinical				
Greenall et al, <sup>26</sup> 1979	1b (RCT)	100 Patients, colon	Clinical AL	Suture bite: 5 mm, AL, 4% (n = 50); 10 mm, 6% (n = 50); not significant
Experimental				
Högström et al, <sup>25</sup> 1985	...	80 Rats	Breaking strength	1.5-mm vs 3.0-mm distance from wound edges; less breaking strength 1.5 mm at day 2
<b>In-between distance of bites</b>				
Experimental				
Waninger et al, <sup>27</sup> 1992	...	432 Rats	Histologic results	2.5-mm vs 1.5-mm suture distance: 1.5 mm, better histologic results
<b>Suture tension</b>				
Experimental				
Waninger et al, <sup>27</sup> 1992	...	432 Rats	ABP, histologic results	No tension vs moderate tension vs high tension on knot: moderate tension, better histologic results; moderate tension, higher ABP at day 4; no tension, higher ABP at day 7
<b>Configuration of the bite<sup>b</sup></b>				
Clinical				
Leslie and Steele, <sup>30</sup> 2003	2b (cohort)	484 Patients; colon	Clinical AL	AL: interrupted serosubmucosal stitches, 0.2%
Pye and Steele, <sup>31</sup> 1996	2b (cohort)	214 Patients; colorectal	Clinical AL	AL: interrupted serosubmucosal stitches, 0.5%
Carty et al, <sup>32</sup> 1991	2b (cohort)	421 Patients; colorectal	Clinical AL	AL: interrupted extramucosal stitches, 2.1%
Lafreniere and Ketcham, <sup>33</sup> 1985	2b (cohort)	134 Patients; colorectal	Clinical AL	AL: interrupted full thickness (modified Gambee), 0
Motson et al, <sup>34</sup> 1984	2b (cohort)	92 Patients, colon	Clinical AL	AL: interrupted full thickness (mattress suture), 4.4%
Experimental				
Krasniqi et al, <sup>29</sup> 2009	...	73 Rats	Histologic results	Serosubmucosal (Halsted) vs full thickness (Gambee) vs posterior Gambee/anterior Halsted stitch: better macroscopic and microscopic histologic results with full thickness
Houdart et al, <sup>28</sup> 1983	...	210 Rats	Histologic results	Extramucosal vs full-thickness stitch: equal histologic results
<b>Inverting vs everting<sup>a</sup></b>				
Clinical				
Goligher et al, <sup>38</sup> 1970	1b (RCT)	70 Patients, colon	Clinical AL	AL: inverting, 2.9% (n = 35); everting, 28.6% (n = 35); significant
Experimental				
Ortiz et al, <sup>35</sup> 1975	...	88 Rats	AL, histologic results	Inverting vs everting: no AL, slower healing, equal adhesions with everting
Irvin and Edwards, <sup>36</sup> 1973	...	93 Rabbits	AL, ABP, histologic results	Inverting 1-layer vs inverting 2-layer vs everting: more AL, lower ABP, delayed mucosal union with everting
Le Douarec and Jouanneau, <sup>37</sup> 1972	...	65 Rabbits	Histologic results	Direct (everting) vs intraluminal (inverting) sutures: direct, more severe inflammation; intraluminal, better histologic repair

Abbreviations: see Table 1.

<sup>a</sup>Highest level of evidence, 1b.<sup>b</sup>Highest level of evidence, 2b.

### In-between Distance of Bites

Lembert described in 1826<sup>125</sup> an in-between distance of approximately 1 cm between sutures. One experimental study conducted by Waninger et al<sup>27</sup> investigating the distance between sutures in rats was included in our review. It concluded that a small distance between sutures (1.5 mm) improves apposition compared with a larger distance (2.5 mm). Neither clinical comparative studies nor cohort studies were found. Again, distances in a rat model are difficult to extrapolate to the patient. Because clinical studies on this topic are lacking, no precise maxim can be distilled from the literature.

### Suture Tension

In routine clinical practice, 2 undefined schools of thought seem to exist: the first believes that sutures should be tightened to prevent dehiscence of the anastomosis, and the sec-

ond considers that sutures should be applied more loosely, allowing maximal perfusion of the cut edges. Only one rat study<sup>27</sup> investigated this, with moderate tension giving the best histologic and microangiographic results. Whether tension on knots could influence the incidence of AL in a clinical setting has not been investigated for interrupted or continuous suturing. On the basis of the literature evaluated in the present review, nothing can be concluded on the proper tension on the thread or the knot.

### Configuration of the Bite

Historically, all opinion leaders proposed their own configuration of gastrointestinal sutures. Anatomic apposition of all layers promoting primary healing was thought to be important. These days, most surgeons use a simple through-all-layers technique. From *ex vivo* studies,<sup>126,127</sup> it is known that sutures through the mucosal layer do not contribute to anastomotic strength.

**Table 3. Included Studies on Single- vs Double-Layer Colorectal Anastomosis<sup>a</sup>**

Source	Level	No. of Cases	Outcome	Results
Everett, <sup>51</sup> 1975	1b (RCT)	92 Patients, colorectal	Clinical AL	AL: 1-layer, 5.0% (n = 40); 2-layer, 4.8% (n = 52); nonsignificant
Ceraldi et al, <sup>52</sup> 1993	2b (nonrandomized)	84 Patients, colon	Clinical AL	AL: 1-layer, 6.8% (n = 44); 2-layer, 9.5% (n = 21); nonsignificant
Fielding et al, <sup>53</sup> 1980	2b (nonrandomized)	1466 Patients, colon	Clinical AL	AL: 1-layer, 12.0% (n = 458); 2-layer, 13.5% (n = 968); nonsignificant
Reichel et al, <sup>47</sup> 1975	2b (nonrandomized)	408 Patients, colorectal	Clinical and radiological leak	AL: 1-layer, 10.3% (n = 320); 2-layer, 10.3% (n = 88); nonsignificant
Athar et al, <sup>48</sup> 1996	...	18 Dogs	Breaking strength	1-Layer, more strength; 2-layer, fewer adhesions, smaller-diameter anastomosis
Langer et al, <sup>46</sup> 1996	...	26 Rabbits	AL, ABP	Corticosteroid model, 1-layer vs 2-layer (vs stapled, NIR): equal AL; 2-layer, higher ABP
Templeton and McKelvey, <sup>42</sup> 1985	...	40 Dogs	AL, ABP, histologic results, collagen	1-Layer vs 2-layer (vs 2× stapled, NIR): AL, 0% vs 30%; histologic results, worse for 2-layer; equal ABP and collagen
Yesilkaya et al, <sup>43</sup> 1985	...	20 Dogs	Collagen	1-Layer vs 2-layer: lower collagen in 2-layer
Graffner et al, <sup>41</sup> 1984	...	18 Pigs	AL, breaking strength, histologic results, blood flow	1-Layer vs 2-layer (vs stapled, NIR): no AL; 2-layer, more inflammation; 2-layer, less blood flow; equal breaking strength
Wheless and Smith, <sup>50</sup> 1983	...	81 Dogs	Blood flow	1-Layer (Gambee) vs 2-layer (vs stapler, NIR): 1-layer, higher blood flow
Schillaci et al, <sup>44</sup> 1979	...	30 Dogs	ABP, collagen	1-Layer vs 2-layer (vs sleeve, NIR): equal ABP and collagen
Chung, <sup>45</sup> 1987	...	30 Dogs	Blood flow	Different hand-sewn vs stapled anastomoses: 2-layer, more reduction of blood flow
Langer, <sup>39</sup> 1975	...	80 Dogs and rats	Histologic results, blood flow	1-Layer vs 2-layer: 2-layer, delayed recovery, more ulceration and stenosing, delayed revascularization
Reichel et al, <sup>47</sup> 1975	...	360 Dogs	AL, ABP, histologic results	1-Layer vs 2-layer: no AL, equal ABP, histologic results worse in 2-layer
Irvin and Edwards, <sup>36</sup> 1973	...	93 Rabbits	ABP, histologic results, collagen	1-Layer vs 2-layer (vs everting, NIR): equal ABP, collagen, and histologic results
Herzog, <sup>49</sup> 1973	...	200 Rats	Breaking strength, blood flow	1-Layer, better vascularization and higher bursting strength
McAdams et al, <sup>40</sup> 1970	...	116 Dogs	Histologic results	1-Layer (Gambee) vs 2-layer (Czerny-Lembert): 2-layer, more inflammation

Abbreviations: ABP, anastomotic bursting pressure; AL, anastomotic leak; NIR, not included in the present review (in case a supplemental subgroup was also studied, outside the topic of this review); RCT, randomized controlled trial; ellipses indicate not applicable.

<sup>a</sup>Highest level of evidence for all studies, 1b.

The present review included 2 experimental studies on rat colon, comparing histologic results of full-thickness sutures with those of serosubmucosal sutures. Houdart et al<sup>28</sup> found no significant histologic differences, but Krasniqi et al<sup>29</sup> found better histologic results for full-thickness sutures with equal anastomotic strength. No comparative clinical studies were found on the configuration of the bite. Because of this lack of evidence, we have included cohort studies,<sup>30-34</sup> reporting low rates of AL for both serosubmucosal and full-thickness suture formats (AL, 0%-4.4%).

We can only conclude, using scarce level 2b evidence from the cohort studies evaluated, that both serosubmucosal and full-thickness sutures seem to provide low rates of AL. It is clear that the configuration of the suture bite is considered of little interest in studies regarding AL.

### Inverting vs Everting Sutures

Since the publication of Lembert,<sup>125</sup> surgeons generally have advocated an inverting technique of gastrointestinal anastomosis because it is believed that protruding mucosa will lead to AL. However, in the 1960s, 2 clinical studies<sup>128,129</sup> showed good healing of everting anastomoses with a low incidence of AL. Between 1960 and 1970,

these 2 noncomparative studies were followed by many experimental publications comparing everting with inverting techniques. They failed to achieve a unanimous conclusion on anastomotic healing; however, they were consistent in showing that everting anastomoses cause more adhesions but less stenosis.<sup>35-37,130-134</sup> All 3 experimental studies<sup>35-37</sup> published after 1970 included in the present review seem to show improved anastomotic healing for inverted anastomoses. The only clinical study matching the inclusion criteria was an RCT<sup>38</sup> showing a 5-fold increased incidence of AL in patients receiving an everting colorectal anastomosis compared with those receiving an inverting colorectal anastomosis. No cohort studies matching our inclusion criteria were found. Therefore, on the basis of available experimental studies and a level 1b clinical study, there seems to be an advantage of inverting over everting colorectal anastomosis; nonetheless, level 1a evidence is lacking.

### SINGLE- VS DOUBLE-LAYER COLORECTAL ANASTOMOSIS

The technique developed by Lembert<sup>125</sup> and later modified by Czerny<sup>135</sup> is based on a double-layer inverting anastomotic technique. In the 19th and the greater part of the 20th centuries, this was the criterion standard for gas-

**Table 4. Included Studies on Interrupted vs Continuous Sutures**

Source	Level <sup>a</sup>	No. of Cases	Outcome	Results
<b>Clinical</b>				
Deen and Smart, <sup>58</sup> 1995	2b (nonrandomized)	53 Patients, colon	Clinical AL	AL: continuous, 3.8% (n = 26); interrupted, 3.8% (n = 27); not significant
Volk et al. <sup>72</sup> 2011	2b (cohort)	463 Patients, ileocolonic	Clinical AL	AL: continuous, 3.1%
Law et al. <sup>66</sup> 1999	2b (cohort)	500 Patients, colorectal	Clinical AL	AL: continuous, 1.4%
AhChong et al. <sup>59</sup> 1996	2b (cohort)	93 Patients, colorectal	Clinical AL	AL: continuous, 2.2%
Flyger et al. <sup>61</sup> 1995	2b (cohort)	105 Patients, colon	Clinical AL	AL: continuous, 1.0%
Max et al. <sup>68</sup> 1991	2b (cohort)	1000 Patients, colorectal	Clinical AL	AL: continuous, 1.0%
Sarin and Lightwood, <sup>70</sup> 1989	2b (cohort)	65 Patients, colon	Clinical AL	AL: continuous, 6.2%
Harder and Vogelbach, <sup>62</sup> 1988	2b (cohort)	143 Patients, colon	Clinical AL	AL: continuous, 0%
Bailey et al. <sup>60</sup> 1984	2b (cohort)	100 Patients, colorectal	Clinical AL	AL: continuous, 0%
Thomson and Robinson, 1993 <sup>71</sup>	2b (cohort)	200 Patients, colorectal	Clinical AL	AL: continuous, 2.0%
Pramateftakis et al. <sup>69</sup> 2010	2b (cohort)	276 Patients, colorectal	Clinical AL	AL: interrupted, 2.5%
Pye and Steele, <sup>31</sup> 1996	2b (cohort)	214 Patients, colorectal	Clinical AL	AL: interrupted, 0.5%
Huguier and Houry, <sup>63</sup> 1982	2b (cohort)	105 Patients, colorectal	Clinical AL	AL: interrupted, 3.8%
Khubchandani and Upson, <sup>65</sup> 1982	2b (cohort)	112 Patients, colorectal	Clinical AL	AL: interrupted, 4.5%
Matheson et al. <sup>67</sup> 1981	2b (cohort)	168 Patients, colorectal	Clinical AL	AL: interrupted, 4.2%
Jonsell and Edelman, <sup>64</sup> 1978	2b (cohort)	165 Patients, colorectal	Clinical AL	AL: interrupted: 8.5%
<b>Experimental</b>				
Shandall et al., <sup>55</sup> 1985	...	40 Rabbits	Blood flow	Continuous vs interrupted: continuous, significantly lower blood flow
Houdart et al., <sup>28</sup> 1983	...	210 Rats	Histologic results	Extramucosal continuous vs extramucosal interrupted vs continuous: equal histologic results at days 2 and 180
Jiborn et al., <sup>73</sup> 1978	...	64 Rats	Collagen	Continuous vs interrupted vs control: continuous, lower collagen metabolism until day 4
Jiborn et al., <sup>57</sup> 1978	...	71 Rats	ABP	Continuous vs interrupted vs control: equal ABP at days 4 and 7
Jiborn et al., <sup>56</sup> 1978	...	71 Rats	Breaking strength	Continuous vs interrupted vs control: equal breaking strength at days 4 and 10
Delaitre et al., <sup>54</sup> 1977	...	83 Rabbits	Histologic results	Continuous vs interrupted: continuous, more evagination of mucus and better apposition (from 1 d to 3 mo)

Abbreviations: ABP, anastomotic bursting pressure; AL, anastomotic leak.  
<sup>a</sup>Highest level of evidence, 2b.

trointestinal anastomosis; in the second half of the 20th century, however, the single-layer anastomosis regained attention through the favorable results obtained by Halstead,<sup>136</sup> Gambee,<sup>137</sup> and Gambee et al.<sup>138</sup> The 13 included experimental studies<sup>36,39-50</sup> came to the same conclusion: double-layer anastomoses are inferior to single-layer anastomoses because of increased inflammation and diminished circulation. One RCT<sup>51</sup> matched the inclusion criteria, showing no significant differences in AL between single- and double-layer colorectal anastomosis in 92 patients. This RCT conducted a subgroup analysis of 25 low colorectal anastomoses, finding a significantly higher incidence of AL in colorectal anastomosis created with the double-layer technique. None of the 3 non-randomized comparative studies<sup>47,52,53</sup> included in this review found a significant difference in AL between the 2 techniques. In conclusion, these results add to the knowledge that single-layer anastomoses take significantly less time to construct and are less costly<sup>139</sup> and are in favor of single anastomoses on the basis of level 1b evidence.

#### INTERRUPTED VS CONTINUOUS SUTURES

The question on whether to use interrupted or continuous sutures arose when single-layer anastomoses became common practice. Six experimental studies were included, showing equivocal results: better serosal apposition<sup>54</sup> and blood flow in continuous sutures,<sup>55</sup> with equal results on ABP and histologic examina-

tion.<sup>28,56,57,73</sup> Randomized controlled trials investigating interrupted and continuous sutures for colorectal anastomosis are lacking; therefore, only 1 small, nonrandomized, comparative clinical study finding no significant differences was included,<sup>58</sup> and noncomparative cohort studies were selected on continuous and interrupted suturing, finding equally low leak rates.<sup>31,59-72</sup> Clinical and experimental studies have not concluded that one technique is superior to the other, and a high level of evidence is lacking (limited here to level 2b); however, from a technical and time-consuming point of view, a continuous suture is preferable over interrupted sutures for creating colorectal anastomosis.

#### HAND-SEWN VS STAPLED COLORECTAL ANASTOMOSIS

After the introduction of stapled colorectal anastomosis in the 1980s, both techniques have become prevalent, without defined indications but for the lower rectal anastomoses. Most surgeons apply both techniques, although often with a personal preference.

Thirteen RCTs<sup>74-86</sup> and 3 meta-analyses<sup>87-89</sup> were included in the present review. Lustosa et al.<sup>87</sup> published a Cochrane meta-analysis of 9 RCTs conducted between 1981 and 1991. In this group of 1233 patients, there was no significant difference in mortality, AL, strictures, or reoperation between stapled and hand-sewn colorectal anastomosis. An earlier meta-analysis,<sup>88</sup> conducted in

**Table 5. Included Studies on Hand-Sewn vs Stapled Colorectal Anastomosis**

Source	Level <sup>a</sup>	No. of Cases	Outcome	Results
<b>Clinical</b>				
Choy et al, <sup>89</sup> 2007	1a (meta-analysis)	955 Patients, ileocolic	Clinical AL	AL: stapled, 1.1% (n = 357); hand-sewn, 3.8% (n = 598); not significant
Lustosa et al, <sup>87</sup> 2001	1a (meta-analysis)	1233 Patients, colorectal	Clinical AL	AL: stapled, 6.3% (n = 622); hand-sewn, 7.1% (n = 611); not significant
MacRae and McLeod, <sup>88</sup> 1998	1a (meta-analysis)	2256 Patients, colorectal	Clinical AL	AL stapled vs hand-sewn: odds ratio, 0.89 (95% CI, 0.58-1.29); not significant
Fingerhut et al, <sup>81</sup> 1995 <sup>b,c</sup>	1b (RCT)	159 Patients, colorectal suprapertitoneal	Clinical AL	AL: stapled, 0 (n = 85); hand-sewn, 0 (n = 74); not significant
Docherty et al, <sup>77</sup> 1995 <sup>c,d</sup>	1b (RCT)	625 Patients, colorectal	Clinical AL	AL: stapled, 4.5% (n = 330); hand-sewn, 4.4% (n = 321); not significant
Fingerhut et al, <sup>80</sup> 1994 <sup>b,c</sup>	1b (RCT)	113 Patients, left infraperitoneal	Clinical AL	AL: stapled, 3.7% (n = 54); hand-sewn, 8.5% (n = 59); not significant
Sarker et al, <sup>86</sup> 1994 <sup>b,c</sup>	1b (RCT)	60 Patients, rectum	Clinical AL	AL: stapled, 0 (n = 30); hand-sewn, 6.7% (n = 30); not significant
Kracht et al, <sup>83</sup> 1993 <sup>b,c,d</sup>	1b (RCT)	268 Patients, colorectal	Clinical AL	AL: stapled, 8.8% (n = 137); hand-sewn, 12.2% (n = 131); not significant
Friend et al, <sup>82</sup> 1990 <sup>c</sup>	1b (RCT)	239 Patients, left colon	Clinical AL	AL: stapled, 3.5% (n = 114); hand-sewn, 8.8% (n = 125); not significant
Cajozzo et al, <sup>76</sup> 1990 <sup>c</sup>	1b (RCT)	48 Patients, colorectal	Clinical AL	AL: stapled, 8.3% (n = 24); hand-sewn, 4.2% (n = 24); not significant
Elhadad, <sup>78</sup> 1990 <sup>b</sup>	1b (RCT)	272 Patients colorectal	Clinical fistula	AL: stapled, 8.3% (n = 139); hand-sewn, 11.5% (n = 133); not significant
Moreno Gonzalez et al, <sup>85</sup> 1989 <sup>b,c</sup>	1b (RCT)	113 Patients, rectum	Clinical AL	AL: stapled, 10.9% (n = 55); hand-sewn, 10.3% (n = 58); not significant
Everett et al, <sup>79</sup> 1986	1b (RCT)	94 Patients, descending colon	Clinical AL	AL: stapled, 0 (n = 44); hand-sewn, 4% (n = 50); not significant
McGinn et al, <sup>84</sup> 1985 <sup>b,c</sup>	1b (RCT)	118 Patients, low colorectal	Clinical AL	AL: stapled, 12.1% (n = 58); hand-sewn, 3.3% (n = 60); significant
Brennan et al, <sup>75</sup> 1982 <sup>c</sup>	1b (RCT)	100 Patients, colorectal	Clinical AL	AL: stapled, 10% (n = 50); hand-sewn, 6% (n = 50); not significant
Beart and Kelly, <sup>74</sup> 1981 <sup>b,c</sup>	1b (RCT)	70 Patients, colorectal	Clinical AL	AL: stapled, 2.9% (n = 35); hand-sewn, 2.9% (n = 35); not significant
Resegotti et al, <sup>99</sup> 2005	2b (nonrandomized)	122 Patients, ileocolic Crohn disease	Clinical AL	AL: stapled, 2.0% (n = 51); hand-sewn, 14.1% (n = 71); significant
Anwar et al, <sup>100</sup> 2004	2b (nonrandomized)	100 Patients, ileocolic malignancy	Clinical AL	AL: stapled, 0 (n = 41); hand-sewn, 0 (n = 59); not significant
Smedh et al, <sup>101</sup> 2002	2b (nonrandomized)	42 Patients, Crohn disease	Clinical AL	AL: STS stapled, 0 (n = 20); ETE hand-sewn, 0 (n = 22); not significant
Sieleznoff et al, <sup>102</sup> 2001	2b (nonrandomized)	116 Patients sigmoid (diverticulitis)	Clinical AL	AL: stapled, 0 (n = 49); hand-sewn, 0 (n = 67); not significant
Montesani et al, <sup>98</sup> 1992	2b (nonrandomized)	533 Patients, colorectal	Clinical AL	AL: stapled, 28.5% (n = 28); hand-sewn, 3.1% (n = 505); significant
Scher et al, <sup>103</sup> 1982	2b (nonrandomized)	242 Patients, colon	Clinical AL	AL: stapled, 2.3% (n = 87); hand-sewn, 2.6% (n = 155); not significant
Adloff et al, <sup>104</sup> 1980	2b (nonrandomized)	51 Patients, rectum	Clinical AL	AL: stapled, 7.7% (n = 26); hand-sewn, 8% (n = 25); not significant
<b>Experimental</b>				
Singer et al, <sup>93</sup> 2004	...	20 Pigs	ABP, histologic results, collagen	Corticosteroid model; equal ABP and collagen; inflammation worse for hand-sewn at day 4
Senagore et al, <sup>92</sup> 1992	...	42 Pigs	ABP, histologic results, blood flow, collagen	Equal ABP, histologic results, blood flow, hydroxyproline
Kent et al, <sup>105</sup> 1992	...	20 Dogs	ABP, histologic results	Hand-sewn: higher ABP; equal histologic results at days 3 and 5
Jansson et al, <sup>90</sup> 1991	...	30 Pigs	AL, breaking strength, blood flow, collagen	Hand-sewn vs stapled (vs glued, NIR): no AL; equal breaking strength, blood flow, and collagen
Dziki et al, <sup>106</sup> 1991	...	24 Dogs	ABP, histologic results, collagen	Hand-sewn: higher ABP at day 4, better histologic results, equal collagen
Julian and Kolachalam, <sup>97</sup> 1989	...	56 Dogs	Blood flow	Equal vascularization at days 3 and 13
Kozol et al, <sup>94</sup> 1988	...	8 Dogs	Histologic results, edema	No significant difference in edema at 28 h
Chung, <sup>45</sup> 1987	...	30 Dogs	Blood flow	Tight stapling: less blood flow; adjusted stapling: better blood flow vs hand-sewn
Graffner et al, <sup>41</sup> 1984	...	18 Pigs	AL, breaking strength, histologic results, blood flow	1-Layer vs 2-layer vs stapled: no AL, more necrosis with stapled, equal blood flow and breaking strength
Moss, <sup>95</sup> 1984	...	10 Dogs	ABP	Stapled, higher ABP at day 4
Buchmann et al, <sup>96</sup> 1983	...	8 Dogs	Histologic results	Equal histologic results at day 4; stapled had more fibrosis at 2, 3, and 6 mo
Wheless and Smith, <sup>50</sup> 1983	...	81 Dogs	Blood flow	1-Layer (Gambee) vs 2-layer vs stapled: stapled had higher blood flow than 1-layer and 2-layer
Polglase et al, <sup>91</sup> 1981	...	24 Dogs	AL, histologic results	Hand-sewn, more narrowing; stapled, more ulcerative gap; equal AL

Abbreviations: see Table 1; ETE, end-to-end; NIR, not included in the present review (in case a supplemental subgroup was also studied, outside the topic of this review); STS, side-to-side.

<sup>a</sup>Highest level of evidence, 1a.

<sup>b</sup>Indicates a randomized controlled trial (RCT) also included in the meta-analysis of Lustosa et al.<sup>87</sup>

<sup>c</sup>Indicates an RCT also included in the meta-analysis of MacRae and McLeod.<sup>88</sup>

<sup>d</sup>Indicates an RCT also included in the meta-analysis of Choy et al.<sup>89</sup>

**Table 6. Included Studies on Hand-Sewn vs Compression Colorectal Anastomosis**

Source	Level <sup>a</sup>	No. of Cases	Outcome	Results
<b>Clinical</b>				
Pählman et al, <sup>116</sup> 1997	1b (RCT)	100 Patients, colon	Clinical AL	AL: BAR, 4% (n = 50); hand-sewn, 2% (n = 50); not significant
Gullichsen et al, <sup>114</sup> 1992	1b (RCT)	150 Patients, colon	Clinical AL	AL: BAR, 2.5% (n = 79); hand-sewn 2-layer, 4.2% (n = 71); not significant
Bubrick et al, <sup>113</sup> 1991	1b (RCT)	782 Patients, colorectal	Clinical AL	AL: BAR, 3% (n = 395); hand-sewn, 3% (n = 283); stapled, 4% (n = 104); not significant
Dyess et al, <sup>115</sup> 1990	1b (RCT)	59 Patients, colon	Clinical AL	AL: BAR, 0 (n = 27); hand-sewn, 0 (n = 16); stapled, 0 (n = 16); not significant
Cahill et al, <sup>111</sup> 1989	1b (RCT)	202 Patients, colorectal	Clinical AL	AL: BAR, 2% (n = 101); hand-sewn, 8.2% (n = 85); stapled, 6.3% (n = 16); not significant
Corman et al, <sup>112</sup> 1989	1b (RCT)	438 Patients, colon	Clinical AL	AL: BAR, 2.7% (n = 222); hand-sewn, 2.5% (n = 162); stapled, 1.9% (n = 54); not significant
<b>Experimental</b>				
Bundy et al, <sup>108</sup> 1993	...	36 Dogs	ABP	BAR vs hand-sewn vs stapled: hand-sewn higher ABP at day 3; equal ABP at day 7
Gullichsen, <sup>110</sup> 1993	...	42 Dogs	Histologic results	BAR vs hand-sewn vs stapled: more edema and inflammation with BAR at days 1 and 7
Smith et al, <sup>107</sup> 1988	...	40 Dogs	AL	2 Sizes BAR vs hand-sewn vs stapled: radiotherapy model; 1.5-mm BAR, more AL; other groups, equal
Maney et al, <sup>109</sup> 1988	...	178 Dogs	AL, ABP, histologic results	BAR vs EEA vs hand-sewn: no AL; equal ABP and histologic results

Abbreviations: see Table 1; BAR, biofragmentable anastomotic ring; EEA, end-to-end anastomosis stapler.

<sup>a</sup>Highest level of evidence, 1b.

**Table 7. Included Studies on Configuration**

Source	Level <sup>a</sup>	No. of Cases	Outcome	Results
<b>Clinical</b>				
Brisinda et al, <sup>119</sup> 2009	1b (RCT)	77 Patients, rectum	Clinical AL	AL: ETE, 29.2% (n = 37); ETS, 5.0% (n = 40); significant
Tsunoda et al, <sup>120</sup> 2009	1b (RCT)	40 Patients, rectum	Clinical AL and stump leak	AL: short (3-cm) limb, 5% (n = 20); long (6-cm) limb, 10% (n = 20); not significant
<b>Experimental</b>				
Willis et al, <sup>118</sup> 2006	...	18 Dogs	Perfusion	Rectum, stapled ETE vs stapled STE (vs J-pouch, NIR): ETE, better blood flow compared with STE
Sailer et al, <sup>117</sup> 2000	...	32 Pigs	Blood flow	Rectum, ETE vs STS (vs small pouch vs large pouch, NIR): equal blood flow

Abbreviations: see Table 1; ETE, end-to-end; ETS, end-to-side; NIR, not included in the present review (in case a supplemental subgroup was also studied, outside the topic of this review); STE, side-to-end; STS, side-to-side.

<sup>a</sup>Highest level of evidence, 1b.

1998, combined 13 RCTs concerning patients with colorectal anastomosis and found similar results: no significant differences in AL or mortality. The Cochrane review conducted by Choy et al<sup>89</sup> included studies on colorectal anastomosis after right hemicolectomy. This review showed significantly less overall AL in the stapled group; however, when clinical AL was used as the only outcome measure, this difference did not reach statistical significance. An interesting subgroup analysis made by Friend et al<sup>82</sup> found more AL in hand-sewn colorectal anastomosis when the anastomoses performed by residents were analyzed separately. Their conclusion was that stapling seems to have an advantage in less-experienced hands. Of 7 included nonrandomized cohort studies included in this review, 5 found no superiority of one technique.<sup>100-104</sup> Two studies found significantly more AL in stapled compared with hand-sewn anastomoses.<sup>98,99</sup> However, one of these had significantly more patients with corticosteroids in the stapled group,<sup>99</sup> while the other included 505 hand-sewn compared with 28 stapled colorectal anastomoses in 20 years.<sup>98</sup> Thirteen experimental studies<sup>41,43,50,90-97,105,106</sup> included herein found results approximately similar to those in the clinical setting: no sig-

nificant differences in AL, with equal or higher ABP in stapled colorectal anastomosis.

In conclusion, the field of hand-sewn vs stapled colorectal anastomosis has been well studied. On the basis of level 1a evidence, no superiority of stapled over hand-sewn colorectal anastomosis exists.

## HAND-SEWN VS COMPRESSION COLORECTAL ANASTOMOSIS

Denans described the first technique to create intestinal anastomoses by compression in 1827,<sup>140</sup> followed by other devices, such as the Murphy button, in 1892.<sup>141</sup> Today the biofragmentable anastomotic ring, made of absorbable polyglycolic acid, is used most often. Four included experimental studies<sup>107-110</sup> showed that compression colorectal anastomosis leads to acceptable healing and strength; 6 included RCTs<sup>111-116</sup> provide equivalent conclusions, finding no significant differences between hand-sewn and compression colorectal anastomosis. Also, noncomparative clinical cohort studies<sup>142-145</sup> including up to 1360 patients have reported incidences of AL between 0.7% and 5%. Although few gastrointestinal sur-

geons routinely use compression colorectal anastomosis, it seems a safe method. On the basis of 6 level 1b studies, no superiority of compression over hand-sewn colorectal anastomosis exists when comparing leak rates.

## CONFIGURATION OF COLORECTAL ANASTOMOSIS

Studies regarding the configuration of the afferent and efferent ileal, colonic, or rectal loops are heterogeneous in patient selection and configuration and often concentrate on stapled pouches for very low anastomoses with outcome variables other than AL. Only 2 experimental studies matched the inclusion criteria; one study<sup>117</sup> found no significant difference in blood flow between ETE or side-to-side anastomosis after rectal resection in pigs, and the other<sup>118</sup> found better blood flow in ETE compared with side-to-end anastomosis after rectal resection in dogs. The included RCTs are also scarce: one<sup>119</sup> on ETE vs end-to-side found more AL in ETE, and the other<sup>120</sup> on the optimum side limb for side-to-end colorectal anastomosis found no significant difference between 3- and 6-cm sized limbs. No studies investigating the ideal length of the enterotomy were identified.

It is difficult to draw a conclusion from this small number of studies; there is one level 1b study showing a lower incidence of AL with end-to-side colorectal anastomosis and one level 1b study indicating that a 3-cm or a 6-cm side limb does not affect the incidence of AL.

## CONCLUSIONS

In the clinical setting, healing of colorectal anastomosis is obscured from direct postoperative inspection. When AL occurs, diagnosis can be made only after the patient has become ill, making it a feared complication with high morbidity and mortality.<sup>1,4,146-148</sup> This systematic review of all aspects of hand-sewn colorectal anastomosis and the comparison of hand-sewn with mechanical anastomosis provides an overview of the existing colorectal anastomotic techniques combined with the available scientific evidence on anastomotic healing. Evaluation of colorectal anastomosis with clinical AL as the outcome measure and proper statistics produced very little level 1 evidence for all aspects of hand-sewn colorectal anastomosis. Nevertheless, we can formulate a conclusion using experimental results combined with clinical results for many aspects: the single-layer continuous suture technique by an inverting technique with slowly absorbable monofilament material seems preferable on the basis of level 1b evidence. However, for the other aspects of the technique, such as how far to place the suture from the anastomotic edge, the intersuture distance in relationship to the distance to the edge, which layers to include in the bite, how high the tension on the suture should be, and through what configuration the anastomosis should be made, surgeons probably rely on their teachers and instinct rather than on scientific evidence.

Large cohort studies that are available, describing low rates of AL for the used anastomotic technique, might indicate that dedicated, high-volume colorectal surgery has a role in lowering the incidence of AL because of a surgeon's familiarity with a certain technique.

Considering mechanical colorectal anastomosis, level 1a evidence indicates that stapling and hand-sewn anastomoses give equal results with regard to clinical AL, and level 1b evidence determines that compression and hand-sewn colorectal anastomosis have similar AL rates. In contrast to all possible variations that exist when sewing an anastomosis by hand, the technique of a stapled technique is much more uniform in the hands of surgeons. This could lead to standardizing colorectal anastomosis and prevent the nonscientific practice of the preferences of individual surgeons from being handed down from teacher to student without documentation of their exact properties and incidence of AL.

We can conclude from this review that, as of now, hand-sewn colorectal anastomosis is constructed following a largely undefined technique. The circumstances of RCTs do not reflect daily practice; therefore, routine, detailed documentation of anastomotic technique of all colorectal operations will be instrumental in formulating a definitive conclusion on the role of the unstandardized hand-sewn colorectal anastomosis.

**Accepted for Publication:** June 25, 2012.

**Correspondence:** Juliette C. Sliker, MD, Department of Surgery, Erasmus Medical Center, Laboratory of Experimental Surgery, Room Ee-173, Postbus 2040, 3000 CA Rotterdam, the Netherlands (juliette\_sliker@hotmail.com).

**Author Contributions:** *Study concept and design:* Sliker, Daams, Jeekel, and Lange. *Acquisition of data:* Sliker and Daams. *Analysis and interpretation of data:* Sliker, Daams, Mulder, Jeekel, and Lange. *Drafting of the manuscript:* Sliker, Daams, and Jeekel. *Critical revision of the manuscript for important intellectual content:* Sliker, Daams, Mulder, Jeekel, and Lange. *Statistical analysis:* Sliker. *Obtained funding:* Jeekel. *Administrative, technical, and material support:* Mulder and Jeekel. *Study supervision:* Jeekel and Lange.

**Conflict of Interest Disclosures:** None reported.

**Online-Only Material:** The eFigure is available at <http://www.jamasurg.com>.

## REFERENCES

- Hyman N, Manchester TL, Osler T, Burns B, Cataldo PA. Anastomotic leaks after intestinal anastomosis: it's later than you think. *Ann Surg*. 2007;245(2):254-258.
- Matthiessen P, Hallböök O, Rutegård J, Simert G, Sjö Dahl R. Defunctioning stoma reduces symptomatic anastomotic leakage after low anterior resection of the rectum for cancer: a randomized multicenter trial. *Ann Surg*. 2007;246(2):207-214.
- Alves A, Panis Y, Trancart D, Regimbeau JM, Pocard M, Valleur P. Factors associated with clinically significant anastomotic leakage after large bowel resection: multivariate analysis of 707 patients. *World J Surg*. 2002;26(4):499-502.
- Peeters KC, Tollenaar RA, Marijnen CA, et al; Dutch Colorectal Cancer Group. Risk factors for anastomotic failure after total mesorectal excision of rectal cancer. *Br J Surg*. 2005;92(2):211-216.
- Komen N, Dijk JW, Lalmahomed Z, et al. After-hours colorectal surgery: a risk factor for anastomotic leakage. *Int J Colorectal Dis*. 2009;24(7):789-795.
- Zorcolo L, Covotta L, Carlomagno N, Bartolo DC. Toward lowering morbidity, mortality, and stoma formation in emergency colorectal surgery: the role of specialization. *Dis Colon Rectum*. 2003;46(11):1461-1468.
- Yeh CY, Changchien CR, Wang JY, et al. Pelvic drainage and other risk factors for leakage after elective anterior resection in rectal cancer patients: a prospective study of 978 patients. *Ann Surg*. 2005;241(1):9-13.
- Matthiessen P, Hallböök O, Andersson M, Rutegård J, Sjö Dahl R. Risk factors for anastomotic leakage after anterior resection of the rectum. *Colorectal Dis*. 2004;6(6):462-469.

9. Rullier E, Laurent C, Garrelon JL, Michel P, Saric J, Parneix M. Risk factors for anastomotic leakage after resection of rectal cancer. *Br J Surg.* 1998;85(3):355-358.
10. Vignali A, Fazio VW, Lavery IC, et al. Factors associated with the occurrence of leaks in stapled rectal anastomoses: a review of 1,014 patients. *J Am Coll Surg.* 1997;185(2):105-113.
11. Mann B, Kleinschmidt S, Stremmel W. Prospective study of hand-sutured anastomosis after colorectal resection. *Br J Surg.* 1996;83(1):29-31.
12. Detry RJ, Kartheuser A, Delriviere L, Saba J, Kestens PJ. Use of the circular stapler in 1000 consecutive colorectal anastomoses: experience of one surgical team. *Surgery.* 1995;117(2):140-145.
13. Hastings JC, Winkle WV, Barker E, Hines D, Nichols W. Effect of suture materials on healing wounds of the stomach and colon. *Surg Gynecol Obstet.* 1975;140(5):701-707.
14. Letwin ER. Evaluation of polyglycolic acid sutures in colon anastomoses. *Can J Surg.* 1975;18(1):30-32.
15. Orringer MB, Appleman HD, Argenta L, Bove E, Cimmino V. Polypropylene suture in esophageal and gastrointestinal operations. *Surg Gynecol Obstet.* 1977;144(1):67-70.
16. Deveney KE, Way LW. Effect of different absorbable sutures on healing of gastrointestinal anastomoses. *Am J Surg.* 1977;133(1):86-94.
17. Foresman PA, Edlich RF, Rodeheaver GT. The effect of new monofilament absorbable sutures on the healing of musculoaponeurotic incisions, gastrotomies, and colonic anastomoses. *Arch Surg.* 1989;124(6):708-710.
18. Lord MG, Broughton AC, Williams HT. A morphologic study on the effect of suturing the submucosa of the large intestine. *Surg Gynecol Obstet.* 1978;146(2):211-216.
19. Munday C, McGinn FP. A comparison of polyglycolic acid and catgut sutures in rat colonic anastomoses. *Br J Surg.* 1976;63(11):870-872.
20. Andersen E, Sondenaa K, Holter J. A comparative study of polydioxanone (PDS) and polyglactin 910 (Vicryl) in colonic anastomoses in rats. *Int J Colorectal Dis.* 1989;4(4):251-254.
21. Pascual I, de Miguel GF, Gómez-Pinedo UA, de Miguel F, Arranz MG, García-Olmo D. Adipose-derived mesenchymal stem cells in biosutures do not improve healing of experimental colonic anastomoses. *Br J Surg.* 2008;95(9):1180-1184.
22. Pasternak B, Rehn M, Andersen L, et al. Doxycycline-coated sutures improve mechanical strength of intestinal anastomoses. *Int J Colorectal Dis.* 2008;23(3):271-276.
23. Clark CG, Wyllie JH, Haggie SJ, Renton P. Comparison of catgut and polyglycolic acid sutures in colonic anastomoses. *World J Surg.* 1977;1(4):501-505.
24. Gillatt DA, Corfield AP, May RE, Bartolo DC, Leaper DJ. Polydioxanone suture in the gastrointestinal tract. *Ann R Coll Surg Engl.* 1987;69(2):54-56.
25. Högstöm H, Haglund U, Zederfeldt B. Suture technique and early breaking strength of intestinal anastomoses and laparotomy wounds. *Acta Chir Scand.* 1985;151(5):441-443.
26. Greenall MJ, Evans M, Pollock AV. Influence of depth of suture bite on integrity of single-layer large-bowel anastomoses: controlled trial. *J R Soc Med.* 1979;72(5):351-356.
27. Waninger J, Kauffmann GW, Shah IA, Farthmann EH. Influence of the distance between interrupted sutures and the tension of sutures on the healing of experimental colonic anastomoses. *Am J Surg.* 1992;163(3):319-323.
28. Houdart R, Lavergne A, Galian A, Hautefeuille P. Anatomic-pathological evolution of single-layer end-to-end digestive anastomoses: a study of 210 colonic anastomoses in rats from the 2nd to the 180th day. *Gastroenterol Clin Biol.* 1983;7(5):465-473.
29. Krasniqi A, Gashi-Luci L, Krasniqi S, et al. A comparison of three single layer anastomotic techniques in the colon of the rat. *Int J Surg.* 2009;7(1):31-35.
30. Leslie A, Steele RJ. The interrupted serosubmucosal anastomosis—still the gold standard. *Colorectal Dis.* 2003;5(4):362-366.
31. Pye G, Steele RJ. Anastomoses involving the colon and rectum: an 8-year experience. *J R Coll Surg Edinb.* 1996;41(2):95-96.
32. Carty NJ, Keating J, Campbell J, Karanjia N, Heald RJ. Prospective audit of an extramucosal technique for intestinal anastomosis. *Br J Surg.* 1991;78(12):1439-1441.
33. Lafreniere R, Ketcham AS. A single layer open anastomosis for all intestinal structures. *Am J Surg.* 1985;149(6):797-798.
34. Motson RW, Bolwell JS, Heath AL, Makin CA, Al Sawaf H. One-layer colonic anastomosis with polyglycolic acid (Dexon) suture: a 3-year prospective audit. *Ann R Coll Surg Engl.* 1984;66(1):19-21.
35. Ortiz H, Azpeitia D, Casalots J, Sitges A. Comparative experimental study of inverting and everting sutures in the colon. *J Chir (Paris).* 1975;109(5-6):691-696.
36. Irvin TT, Edwards JP. Comparison of single-layer inverting, two-layer inverting, and everting anastomoses in the rabbit colon. *Br J Surg.* 1973;60(6):453-457.
37. Le Douarec P, Jouanneau P. Colic anastomosis: experimental study of sutures in one plane of the rabbit colon; comparison of direct and intraluminal sutures. *J Chir (Paris).* 1972;104(5):451-464.
38. Goligher JC, Morris C, McAdam WA, De Dombal FT, Johnston D. A controlled trial of inverting versus everting intestinal suture in clinical large-bowel surgery. *Br J Surg.* 1970;57(11):817-822.
39. Langer S. Complex investigation of the efficiency of large bowel anastomosis techniques (clinical and experimental studies). *Chirurgia Gastroenterologica.* 1975;9:69-80.
40. McAdams AJ, Meikle AG, Taylor JO. One layer or two layer colonic anastomoses? *Am J Surg.* 1970;120(4):546-550.
41. Graffner H, Andersson L, Löwenhielm P, Walther B. The healing process of anastomoses of the colon: a comparative study using single, double-layer or stapled anastomosis. *Dis Colon Rectum.* 1984;27(12):767-771.
42. Templeton JL, McKelvey ST. Low colorectal anastomoses: an experimental assessment of two sutured and two stapled techniques. *Dis Colon Rectum.* 1985;28(1):38-41.
43. Yesilkaya Y, Soyhan N, Bengisu N, Sen M, Aritas Y. The effects of different suture techniques on collagen metabolism in experimental distal colonic anastomoses. *Br J Surg.* 1985;72(12):987-989.
44. Schillaci A, Cavallaro A, Stipa S. Comparative results of three different techniques for colonic anastomosis in the dog. *Surg Gynecol Obstet.* 1979;149(2):238-240.
45. Chung RS. Blood flow in colonic anastomoses: effect of stapling and suturing. *Ann Surg.* 1987;206(3):335-339.
46. Langer JC, Srinathan SK, Pelletier GJ. Effect of surgical technique on intestinal anastomotic healing in steroid-treated rabbits. *Dig Surg.* 1996;13(3):205-208.
47. Reichel K, Rauner P, Guthy E. Clinical and experimental evaluation of single and double layer entero anastomosis. *Chirurgia Gastroenterologica.* 1975;9:461-467.
48. Athar M, Chaudhry NI, Shakoor A, Khan MA. Studies on end-to-end colonic anastomosis in the dog: a comparison of techniques. *Acta Vet Hung.* 1996;44(3):349-356.
49. Herzog B. The one-layer and two-layer intestinal anastomosis in animal experiments. *Prog Pediatr Surg.* 1973;5:37-59.
50. Wheelless CR Jr, Smith JJ. A comparison of the flow of iodine 125 through three different intestinal anastomoses: standard, Gambee, and stapler. *Obstet Gynecol.* 1983;62(4):513-518.
51. Everett WG. A comparison of one layer and two layer techniques for colorectal anastomosis. *Br J Surg.* 1975;62(2):135-140.
52. Ceraldi CM, Rypins EB, Monahan M, Chang B, Sarfeh IJ. Comparison of continuous single layer polypropylene anastomosis with double layer and stapled anastomoses in elective colon resections. *Am Surg.* 1993;59(3):168-171.
53. Fielding LP, Stewart-Brown S, Blesovsky L, Kearney G. Anastomotic integrity after operations for large-bowel cancer: a multicentre study. *Br Med J.* 1980;281(6237):411-414.
54. Delaitre B, Champault G, Chapuis Y, Patel JC, Louvel A, Leger L. Continuous and interrupted intestinal sutures: experimental and clinical study [in French]. *J Chir (Paris).* 1977;113(1):43-57.
55. Shandall A, Lowndes R, Young HL. Colonic anastomotic healing and oxygen tension. *Br J Surg.* 1985;72(8):606-609.
56. Jiborn H, Ahonen J, Zederfeldt B. Healing of experimental colonic anastomoses, II: breaking strength of the colon after left colon resection and anastomosis. *Am J Surg.* 1978;136(5):595-599.
57. Jiborn H, Ahonen J, Zederfeldt B. Healing of experimental colonic anastomoses, I: bursting strength of the colon after left colon resection and anastomosis. *Am J Surg.* 1978;136(5):587-594.
58. Deen KI, Smart PJ. Prospective evaluation of sutured, continuous, and interrupted single layer colonic anastomoses. *Eur J Surg.* 1995;161(10):751-753.
59. AhChong AK, Chiu KM, Law IC, Chu MK, Yip AW. Single-layer continuous anastomosis in gastrointestinal surgery: a prospective audit. *Aust N Z J Surg.* 1996;66(1):34-36.
60. Bailey HR, LaVoo JW, Max E, Smith KW, Butts DR, Hampton JM. Single-layer polypropylene colorectal anastomosis: experience with 100 cases. *Dis Colon Rectum.* 1984;27(1):19-23.
61. Flyger HL, Håkansson TU, Jensen LP. Single layer colonic anastomosis with a continuous absorbable monofilament polyglyconate suture. *Eur J Surg.* 1995;161(12):911-913.
62. Harder F, Vogelbach P. Single-layer end-on continuous suture of colonic anastomoses. *Am J Surg.* 1988;155(4):611-614.

63. Huguier M, Houry S. Manual colorectal anastomosis: immediate results [in French]. *Nouv Presse Med*. 1982;11(29):2211-2213.
64. Jonsell G, Edelmann G. Single-layer anastomosis of the colon: a review of 165 cases. *Am J Surg*. 1978;135(5):630-632.
65. Khubchandani M, Upson JF. Single-layer anastomosis of the colon and rectum. *Dis Colon Rectum*. 1982;25(2):113-117.
66. Law WL, Bailey HR, Max E, et al. Single-layer continuous colon and rectal anastomosis using monofilament absorbable suture (Maxon): study of 500 cases. *Dis Colon Rectum*. 1999;42(6):736-740.
67. Matheson NA, Valerio D, Farquharson A, Thomson H. Single-layer anastomosis in the large bowel: ten years' experience. *J R Soc Med*. 1981;74(1):44-48.
68. Max E, Sweeney WB, Bailey HR, et al. Results of 1,000 single-layer continuous polypropylene intestinal anastomoses. *Am J Surg*. 1991;162(5):461-467.
69. Pramateftakis MG, Vrakas G, Hatzigianni P, et al. The handsewn anastomosis after colon resection due to colonic cancer. *Tech Coloproctol*. 2010;14(suppl 1):S57-S59.
70. Sarin S, Lightwood RG. Continuous single-layer gastrointestinal anastomosis: a prospective audit. *Br J Surg*. 1989;76(5):493-495.
71. Thomson WH, Robinson MH. One-layer continuously sutured colonic anastomosis. *Br J Surg*. 1993;80(11):1450-1451.
72. Volk A, Kersting S, Held HC, Saeger HD. Risk factors for morbidity and mortality after single-layer continuous suture for ileocolonic anastomosis. *Int J Colorectal Dis*. 2011;26(3):321-327.
73. Jiborn H, Ahonen J, Zederfeldt B. Healing of experimental colonic anastomoses: the effect of suture technic on collagen concentration in the colonic wall. *Am J Surg*. 1978;135(3):333-340.
74. Beart RW Jr, Kelly KA. Randomized prospective evaluation of the EEA stapler for colorectal anastomoses. *Am J Surg*. 1981;141(1):143-147.
75. Brennan SS, Pickford IR, Evans M, Pollock AV. Staples or sutures for colonic anastomoses—a controlled clinical trial. *Br J Surg*. 1982;69(12):722-724.
76. Cajozzo M, Compagno G, DiTora P, Spallitta SI, Bazan P. Advantages and disadvantages of mechanical vs. manual anastomosis in colorectal surgery: a prospective study. *Acta Chir Scand*. 1990;156(2):167-169.
77. Docherty JG, McGregor JR, Akyol AM, Murray GD, Galloway DJ, West of Scotland and Highland Anastomosis Study Group. Comparison of manually constructed and stapled anastomoses in colorectal surgery. *Ann Surg*. 1995;221(2):176-184.
78. Elhadad A. Colorectal anastomosis: manual or mechanical? a controlled multicenter study [in French]. *Chirurgie*. 1990;116(4-5):425-428.
79. Everett WG, Friend PJ, Forty J. Comparison of stapling and hand-suture for left-sided large bowel anastomosis. *Br J Surg*. 1986;73(5):345-348.
80. Fingerhut A, Elhadad A, Hay JM, Lacaine F, Flamant Y; French Associations for Surgical Research. Infraperitoneal colorectal anastomosis: hand-sewn versus circular staples—a controlled clinical trial. *Surgery*. 1994;116(3):484-490.
81. Fingerhut A, Hay JM, Elhadad A, Lacaine F, Flamant Y; French Associations for Surgical Research. Supraperitoneal colorectal anastomosis: hand-sewn versus circular staples—a controlled clinical trial. *Surgery*. 1995;118(3):479-485.
82. Friend PJ, Scott R, Everett WG, Scott IH. Stapling or suturing for anastomoses of the left side of the large intestine. *Surg Gynecol Obstet*. 1990;171(5):373-376.
83. Kracht M, Hay JM, Fagniez PL, Fingerhut A. Ileocolonic anastomosis after right hemicolectomy for carcinoma: stapled or hand-sewn? a prospective, multicenter, randomized trial. *Int J Colorectal Dis*. 1993;8(1):29-33.
84. McGinn FP, Gartell PC, Clifford PC, Brunton FJ. Staples or sutures for low colorectal anastomoses: a prospective randomized trial. *Br J Surg*. 1985;72(8):603-605.
85. Moreno Gonzalez E, Rico Selas P, Mansilla Molina D, et al. Results of surgery for cancer of the rectum with sphincter conservation: a randomized study on instrumental versus manual anastomosis. *Acta Oncol*. 1989;28(2):241-244.
86. Sarker SK, Chaudhry R, Sinha VK. A comparison of stapled vs handsewn anastomosis in anterior resection for carcinoma rectum. *Indian J Cancer*. 1994;31(2):133-137.
87. Lustosa SA, Matos D, Atallah AN, Castro AA. Stapled versus handsewn methods for colorectal anastomosis surgery. *Cochrane Database Syst Rev*. 2001;(3):CD003144.
88. MacRae HM, McLeod RS. Handsewn vs. stapled anastomoses in colon and rectal surgery: a meta-analysis. *Dis Colon Rectum*. 1998;41(2):180-189.
89. Choy PY, Bissett IP, Docherty JG, Parry BR, Merrie AE. Stapled versus hand-sewn methods for ileocolic anastomoses. *Cochrane Database Syst Rev*. 2007;(3):CD004320.
90. Jansson OK, Zilling TL, Walther BS. Healing of colonic anastomoses: comparative experimental study of glued, manually sutured, and stapled anastomoses. *Dis Colon Rectum*. 1991;34(7):557-562.
91. Polglase AL, Hughes ES, McDermott FT, Pihl E, Burke FR. A comparison of end-to-end staple and suture colorectal anastomosis in the dog. *Surg Gynecol Obstet*. 1981;152(6):792-796.
92. Senagore A, Milsom JW, Walshaw RK, Dunston R, Chaudry IH. Direct comparison between Czerny-Lembert and circular-stapled anastomotic techniques in colorectal anastomosis: a similar pattern of healing for both. *Dis Colon Rectum*. 1992;35(9):862-869.
93. Singer MA, Cintron JR, Benedetti E, Lamba A, Abcarian H. Hand-sewn versus stapled intestinal anastomoses in a chronically steroid-treated porcine model. *Am Surg*. 2004;70(2):151-156.
94. Kozol RA, Mulligan M, Downes RJ, Forouhar FA, Kreutzer DL. Early colonic anastomotic edema: evaluation of stapled vs. hand-sewn anastomoses. *Dis Colon Rectum*. 1988;31(7):503-506.
95. Moss G. Colorectal anastomotic strength: staples vs conventional sutures. *J Abdom Surg*. 1984;26:73-77.
96. Buchmann P, Schneider K, Gebbers JO. Fibrosis of experimental colonic anastomosis in dogs after EEA stapling or suturing. *Dis Colon Rectum*. 1983;26(4):217-220.
97. Julian TB, Kolachalam RB. Microangiographic study of healing wounds in canine intestinal anastomoses. *Vasc Surg*. 1989;23(4):296-303.
98. Montesani C, De Milito R, Chiappalone S, Narilli P, D'Amato A, Ribotta G. Critical evaluation of the anastomoses in large bowel surgery: experience gained in 533 cases. *Hepatogastroenterology*. 1992;39(4):304-308.
99. Resegotti A, Astegiano M, Farina EC, et al. Side-to-side stapled anastomosis strongly reduces anastomotic leak rates in Crohn's disease surgery. *Dis Colon Rectum*. 2005;48(3):464-468.
100. Anwar S, Hughes S, Eadie AJ, Scott NA. Anastomotic technique and survival after right hemicolectomy for colorectal cancer. *Surgeon*. 2004;2(5):277-280.
101. Smedh K, Andersson M, Johansson H, Hagberg T. Preoperative management is more important than choice of sutured or stapled anastomosis in Crohn's disease. *Eur J Surg*. 2002;168(3):154-157.
102. Sieleznoff I, Malouf AJ, Pirro N, Cesari J, Brunet C, Sastre B. Short-term functional outcome following elective surgery for complicated sigmoid diverticular disease: sutured or stapled end-to-end anastomosis to the proximal rectum? *Colorectal Dis*. 2001;3(1):23-27.
103. Scher KS, Scott-Conner C, Jones CW, Leach M. A comparison of stapled and sutured anastomoses in colonic operations. *Surg Gynecol Obstet*. 1982;155(4):489-493.
104. Adloff M, Arnaud JP, Beehary S. Stapled vs sutured colorectal anastomosis. *Arch Surg*. 1980;115(12):1436-1438.
105. Kent C, Warner K, Miller J, Schreiber H. Ileocolonic anastomosis: a comparison of the patency of stapled versus hand-sewn techniques. *Am Surg*. 1992;58(10):638-640.
106. Dziki AJ, Duncan MD, Harmon JW, et al. Advantages of handsewn over stapled bowel anastomosis. *Dis Colon Rectum*. 1991;34(6):442-448.
107. Smith AD, Bubrick MP, Mestitz ST, et al. Evaluation of the biofragmentable anastomotic ring following preoperative irradiation to the rectosigmoid in dogs. *Dis Colon Rectum*. 1988;31(1):5-9.
108. Bundy CA, Jacobs DM, Zera RT, Bubrick MP. Comparison of bursting pressure of sutured, stapled and BAR anastomoses. *Int J Colorectal Dis*. 1993;8(1):1-3.
109. Maney JW, Katz AR, Li LK, Pace WG, Hardy TG. Biofragmentable bowel anastomosis ring: comparative efficacy studies in dogs. *Surgery*. 1988;103(1):56-62.
110. Gullichsen R. The biofragmentable ring in intestinal surgery. *Eur J Surg Suppl*. 1993;(569):1-31.
111. Cahill CJ, Betzler M, Gruwez JA, Jeekel J, Patel JC, Zederfeldt B. Sutureless large bowel anastomosis: European experience with the biofragmentable anastomosis ring. *Br J Surg*. 1989;76(4):344-347.
112. Corman ML, Prager ED, Hardy TG Jr, Bubrick MP. Comparison of the Valtrac biofragmentable anastomosis ring with conventional suture and stapled anastomosis in colon surgery: results of a prospective, randomized clinical trial. *Dis Colon Rectum*. 1989;32(3):183-187.
113. Bubrick MP, Corman ML, Cahill CJ, Hardy TG Jr, Nance FC, Shatney CH; BAR Investigational Group. Prospective, randomized trial of the biofragmentable anastomosis ring. *Am J Surg*. 1991;161(1):136-143.
114. Gullichsen R, Havia T, Ovaska J, Rantala A. Colonic anastomosis using the biofragmentable anastomotic ring and manual suture: a prospective, randomized study. *Br J Surg*. 1992;79(6):578-580.
115. Dyess DL, Curreri PW, Ferrara JJ. A new technique for sutureless intestinal anastomosis: a prospective, randomized, clinical trial. *Am Surg*. 1990;56(2):71-75.
116. Pahlman L, Ejerblad S, Graf W, et al. Randomized trial of a biofragmentable bowel anastomosis ring in high-risk colonic resection. *Br J Surg*. 1997;84(9):1291-1294.

117. Sailer M, Debus ES, Fuchs KH, Beyerlein J, Thiede A. Comparison of anastomotic microcirculation in coloanal J-pouches versus straight and side-to-end coloanal reconstruction: an experimental study in the pig. *Int J Colorectal Dis.* 2000;15(2):114-117.
118. Willis S, Hölzl F, Krones CJ, Tittel A, Schumpelick V. Evaluation of anastomotic microcirculation after low anterior rectal resection: an experimental study with different reconstruction forms in dogs. *Tech Coloproctol.* 2006;10(3):222-226.
119. Brisinda G, Vanella S, Cadeddu F, et al. End-to-end versus end-to-side stapled anastomoses after anterior resection for rectal cancer. *J Surg Oncol.* 2009;99(1):75-79.
120. Tsunoda A, Kamiyama G, Narita K, Watanabe M, Nakao K, Kusano M. Prospective randomized trial for determination of optimum size of side limb in low anterior resection with side-to-end anastomosis for rectal carcinoma. *Dis Colon Rectum.* 2009;52(9):1572-1577.
121. Durdley P, Bucknall TE. Assessment of sutures for use in colonic surgery: an experimental study. *J R Soc Med.* 1984;77(6):472-477.
122. Houdart R, Lavergne A, Valleur P, Hautefeuille P. Polydioxanone in digestive surgery: an experimental study. *Am J Surg.* 1986;152(3):268-271.
123. Lerwick E. Studies on the efficacy and safety of polydioxanone monofilament absorbable suture. *Surg Gynecol Obstet.* 1983;156(1):51-55.
124. Ray JA, Doddi N, Regula D, Williams JA, Melveger A. Polydioxanone (PDS), a novel monofilament synthetic absorbable suture. *Surg Gynecol Obstet.* 1981;153(4):497-507.
125. Lambert A. Mémoire sur L'entéroraphie avec la description d'un procédé nouveau pour pratiquer cette opération chirurgicale. In: Breschet MG, ed. *Répertoire Général d'Anatomie et de Physiologie Pathologique et des Cliniques Chirurgicales.* Vol 2. Paris, France: Boiste Fils; 1826:100-107.
126. Egorov VI, Schastlivtsev V, Turusov RA, Baranov AO. Participation of the intestinal layers in supplying of the mechanical strength of the intact and sutured gut. *Eur Surg Res.* 2002;34(6):425-431.
127. Tera H, Aberg C. Tissue holding power to a single suture in different parts of the alimentary tract. *Acta Chir Scand.* 1976;142(5):343-348.
128. Buyers RA, Meier LA. Everting suture of the bowel: experimental and clinical experience in duodenal closure and colorectal anastomosis. *Surgery.* 1968;63(3):475-480.
129. Getzen LC. Clinical use of everted intestinal anastomoses. *Surg Gynecol Obstet.* 1966;123(5):1027-1036.
130. Ballantyne GH. The experimental basis of intestinal suturing: effect of surgical technique, inflammation, and infection on enteric wound healing. *Dis Colon Rectum.* 1984;27(1):61-71.
131. Thornton FJ, Barbul A. Healing in the gastrointestinal tract. *Surg Clin North Am.* 1997;77(3):549-573.
132. Yale CE, Van Gemert JV. Healing of inverted and everted intestinal anastomoses in germfree rats. *Surgery.* 1971;69(3):382-388.
133. Getzen LC, Roe RD, Holloway CK. Comparative study of intestinal anastomotic healing in inverted and everted closures. *Surg Gynecol Obstet.* 1966;123(6):1219-1227.
134. Garner A, Hargreaves AW, Keddle NC. Colonic anastomosis: a histopathological study in the rabbit. *Br J Surg.* 1969;56(9):673-676.
135. Czerny V. Zur Darmresektion. *Berl Klin Wschr.* 1880;17:637.
136. Halstead W. Circular suture of the intestine: an experimental study. *Am J Med Sci.* 1887;94:436-461.
137. Gambee LP. A single-layer open intestinal anastomosis applicable to the small as well as the large intestine. *West J Surg Obstet Gynecol.* 1951;59(1):1-5.
138. Gambee LP, Garnjobst W, Hardwick CE. Ten years' experience with a single layer anastomosis in colon surgery. *Am J Surg.* 1956;92(2):222-227.
139. Burch JM, Franciose RJ, Moore EE, Biffi WL, Offner PJ. Single-layer continuous versus two-layer interrupted intestinal anastomosis: a prospective randomized trial. *Ann Surg.* 2000;231(6):832-837.
140. Denans F. *Nouveau procédé pour la guérison des plaies des intestins: recueil de la société Royale de Médecine de Marseille.* Marseille, France: Imprimerie d'Archard; 1827:4.
141. Murphy JB. Cholecysto-intestinal, gastro-intestinal, entero-intestinal anastomosis, and approximation without sutures. *Medical Record.* 1892;42:665-676.
142. Wullstein C, Gross E. Compression anastomosis (AKA-2) in colorectal surgery: results in 442 consecutive patients. *Br J Surg.* 2000;87(8):1071-1075.
143. Kim SH, Choi HJ, Park KJ, et al. Sutureless intestinal anastomosis with the biofragmentable anastomosis ring: experience of 632 anastomoses in a single institute. *Dis Colon Rectum.* 2005;48(11):2127-2132.
144. Di Castro A, Biancari F, Brocato R, Adami EA, Truosolo B, Massi G. Intestinal anastomosis with the biofragmentable anastomosis ring. *Am J Surg.* 1998;176(5):472-474.
145. Thiede A, Geiger D, Dietz UA, et al. Overview on compression anastomoses: biofragmentable anastomosis ring multicenter prospective trial of 1666 anastomoses. *World J Surg.* 1998;22(1):78-87.
146. Alberts JC, Parvaiz A, Moran BJ. Predicting risk and diminishing the consequences of anastomotic dehiscence following rectal resection. *Colorectal Dis.* 2003;5(5):478-482.
147. Branagan G, Finnis D; Wessex Colorectal Cancer Audit Working Group. Prognosis after anastomotic leakage in colorectal surgery. *Dis Colon Rectum.* 2005;48(5):1021-1026.
148. Fielding LP, Phillips RK, Hittinger R. Factors influencing mortality after curative resection for large bowel cancer in elderly patients. *Lancet.* 1989;1(8638):595-597.