

Original Investigation

Ultrasonography-Guided Bilateral Rectus Sheath Block vs Local Anesthetic Infiltration After Pediatric Umbilical Hernia Repair

A Prospective Randomized Clinical Trial

R. Scott Dingeman, MD; Lindsay M. Barus, MPH; Hyun Kee Chung, MD; David J. Clendenin, MD; Christopher S. Lee, MD; Sarah Tracy, BA; Victor M. Johnson, MA; Kate V. Dennett, BA; David Zurakowski, PhD; Catherine Chen, MD, MPH

← Invited Commentary page 713

IMPORTANCE Regional anesthetic techniques can be used to alleviate postoperative pain in children undergoing pediatric surgical procedures. Use of ultrasonographic guidance for bilateral rectus sheath block (BRSB) has been shown to improve immediate pain scores and reduce use of postoperative analgesia in the postanesthesia care unit (PACU).

OBJECTIVE To compare efficacy of ultrasonography-guided BRSB and local anesthetic infiltration (LAI) in providing postoperative analgesia after pediatric umbilical hernia repair.

DESIGN Prospective, observer-blinded, randomized clinical trial.

SETTING Tertiary-referral urban children's hospital.

PARTICIPANTS Eligible children 3 to 12 years of age undergoing elective umbilical hernia repair from November 16, 2009, through May 31, 2011.

INTERVENTIONS Ropivacaine hydrochloride administered at the conclusion of surgery as LAI by the surgeon (n = 25) or as ultrasonography-guided BRSB by the anesthesiologist (n = 27).

MAIN OUTCOMES AND MEASURES Scores on the FACES Pain Rating Scale measured at 10-minute intervals and all use of analgesic medications in the PACU.

RESULTS Median FACES scores in the PACU were lower in the BRSB group compared with the LAI group at 10 minutes (0 vs 1; $P = .04$), 30 minutes (0 vs 1; $P = .01$), and 40 minutes or later (0 vs 1; $P = .03$). Fewer doses of opioid and nonopioid medications were given to the BRSB group compared with the LAI group (5 vs 11 doses for opioids; 5 vs 10 for nonopioids).

CONCLUSIONS AND RELEVANCE In the PACU, ultrasonography-guided BRSB after umbilical hernia repair in children is associated with lower median FACES scores and decreased use of opioid and nonopioid medications compared with LAI. Future studies could examine the use of longer-acting anesthetic agents with ultrasonography-guided BRSB.

TRIAL REGISTRATION clinicaltrials.gov Identifier: NCT01015053

JAMA Surg. 2013;148(8):707-713. doi:10.1001/jamasurg.2013.1442
Published online June 12, 2013.

Author Affiliations: Author affiliations are listed at the end of this article.

Corresponding Author: Catherine Chen, MD, MPH, Department of Surgery, Boston Children's Hospital, 300 Longwood Ave, Boston, MA 02115 (catherine.chen@childrens.harvard.edu).

Regional anesthetic techniques have been used to alleviate postoperative pain in children undergoing outpatient surgical procedures since the mid-1990s. Bilateral rectus sheath block (BRSB) was initially described for use in pediatric patients undergoing umbilical hernia repair without ultrasonographic guidance,^{1,2} with 1 randomized pilot study demonstrating no advantage of BRSB over local anesthetic infiltration (LAI) for postoperative pain management.³ Use of ultrasonographic guidance for ilioinguinal and/or iliohypogastric nerve blocks in children was subsequently shown to reduce the volume of the local anesthetic used.^{4,5} This technical advancement was then used for BRSB to enhance identification of rectus sheath anatomy⁶ and was shown in a small study of 20 children undergoing elective umbilical hernia repair to provide sufficient pain control without the need for additional postoperative analgesia.⁷ Gurnaney et al⁸ recently conducted a randomized prospective study to compare ultrasonography-guided BRSB and LAI in providing postoperative analgesia after pediatric umbilical hernia repair. Although the timing of the placement of the local anesthetic was different between the 2 groups, they found that BRSB decreased the amount of opioid pain medication used during the perioperative period.

Before our study, the accepted regimen for postoperative pain management after elective umbilical hernia repair at our institution consisted of surgeon-administered LAI in the wound at the end of surgery followed by a combination of opioid and nonopioid pain medications in the postanesthesia care unit (PACU) and at home. We undertook this prospective, randomized, observer-blinded study to compare the efficacy of ultra-

sonography-guided BRSB and LAI in providing self-reported postoperative analgesia after umbilical hernia repair in children. Our primary objective was to compare patient-reported pain scores using the FACES Pain Rating Scale (FACES) developed by Wong and Baker⁹ at defined intervals in the PACU and at home within the first 24 hours after surgery. Our secondary aims were to compare the use of opioid and nonopioid medications between treatment groups in the PACU and at home and to obtain measures of parents' perceptions of their child's pain at 12 and 24 hours after surgery.

Methods

The study design was a prospective, randomized, observer-blinded trial with 2 parallel arms. Approval was obtained from the institutional review board of Boston Children's Hospital before patient enrollment (09-08-0398). Consent was obtained from each patient's legal guardian.

Participants

Children 3 to 12 years of age who underwent elective umbilical hernia repair from November 16, 2009, through May 31, 2011, at our institution were eligible for this study (Figure 1). Exclusion criteria consisted of the following: American Society of Anesthesiologists classification of III or greater, history of a complex regional pain syndrome, history of long-term analgesic use, use of any analgesic (eg, an opioid medication, acetaminophen, or a nonsteroidal anti-inflammatory agent) within 24 hours before surgery, history of renal insufficiency or a bleeding disorder, concurrent additional surgery at another anatomic site, being a ward of the state, a non-English- or non-Spanish-speaking patient or primary caregiver, inability to document postoperative pain level using FACES scores, and inability of the primary caregiver to comply with home instructions.


Randomization and Treatment Allocation

Randomization proceeded after written informed consent was obtained in the preoperative area and was assigned by a research team member (L.M.B.) blinded to the group allocation. The randomization scheme was created using a uniform (0 or 1) random number generator in commercially available statistical software (SAS, version 9.2; SAS Institute, Inc) incorporating an age-stratified permuted block method of size 4 to achieve optimal balance (patient age strata, 3-7 and 8-12 years). Sealed envelopes for each age stratum containing the random allocations were opaque and tamperproof.

Blinding

The surgeon (including C.C.), regional pediatric anesthesiologist (R.S.D., H.K.C., D.J.C., and C.S.L.), and operating room staff (anesthesiologists and nurses) were not blinded to the treatment group. However, the patient and family, recovery room nurses, and study coordinator who collected the data from families were blinded. All patients received 2 uniform sterile dressings that were applied to the surgical site as well as the potential regional block sites regardless of randomization.

Figure 1. CONSORT Flow Diagram



This diagram uses the criteria of 133 patients with umbilical hernia aged 3 to 12 years who underwent surgical repair at Boston Children's Hospital from November 16, 2009, through May 31, 2011. BRSB indicates bilateral rectus sheath block; LAI, local anesthetic infiltration.

Interventions

Thirteen staff surgeons enrolled their patients in the study and performed umbilical hernia repairs using standard umbilical incisions. Four pediatric anesthesiologists with expertise in regional anesthesia performed the BRSB. Patients were randomized to one of the following 2 treatment arms using equivalent, volume-adjusted doses of ropivacaine hydrochloride: (1) LAI injected at the site of incision by the surgeon or (2) ultrasonography-guided BRSB administered by the pediatric regional anesthesiologist at the conclusion of surgery.

Study Protocol

Anesthetic Technique

Patients received no acetaminophen or midazolam before surgery. Anesthesia was induced via a mask using inhalational anesthetics and oxygen. After peripheral intravenous (IV) placement, general anesthesia was maintained via laryngeal mask airway or endotracheal tube depending on the intraoperative anesthesiologist's preference. Intravenous fentanyl citrate (1 µg/kg) was administered before incision for intraoperative analgesia. All study patients received IV ketorolac tromethamine, 0.5 mg/kg, at the end of the case. Before emergence, patients received IV ondansetron hydrochloride, 0.1 mg/kg.

Description of Treatment Arms

At the conclusion of surgery, patients in the BRSB underwent cleansing of the periumbilical abdominal wall beyond the lateral border of the rectus muscle with the external oblique muscle bilaterally using a 2% concentration of chlorhexidine gluconate. A ultrasonographic probe (M-Turbo with the L25x transducer; SonoSite, Inc) was used to image the lateral border of the rectus sheath cephalad to the level of the umbilicus. A 22-g short-bevel needle was used to introduce 0.2% ropivacaine hydrochloride, 0.5 mL/kg, between the posterior rectus sheath and the rectus muscle with spread of the anesthetic agent monitored by ultrasonography guidance. This procedure was repeated on the opposite side.

For the LAI arm, patients received a subcutaneous and/or intradermal injection of 0.5% ropivacaine hydrochloride, 0.4 mL/kg, at the surgical site at the conclusion of the surgical procedure.

Data Collection

Demographics and clinical data were collected from patients' medical records. Race was obtained from hospital records and was defined by the patient. Preoperative vital signs and a baseline FACES score were documented in the preoperative area before induction of anesthesia. Surgical start and stop times (which includes time required for LAI by the surgeon or BRSB by the anesthesiologist) were documented.

A blinded research team member assessed postoperative pain using FACES scores once the patient was awake and able to respond to questions and continuing at 10-minute intervals. Analgesics in the PACU were administered according to the following patient-reported FACES scores: no medication for scores of 0 or 1; oral acetaminophen, 12.5 mg/kg, for scores of 2 or 3; and IV morphine sulfate, 0.05 mg/kg (up to a maxi-

um of 0.1 mg/kg), or oral codeine phosphate, 1 mg/kg (if tolerating oral intake), for scores of 4 or 5.

Caregivers recorded data on discharge in a study journal. The FACES scores were documented at home at 4-hour intervals after discharge from the PACU and up to 24 hours after discharge. Oral analgesics were administered and documented according to the patient's self-reported FACES scores as described for the PACU protocol. Sleeping children were not awakened because they were assumed to be resting comfortably. All other medications (natural or chemical) given during the 24-hour home period were also documented. Caregivers also completed the Parent's Postoperative Pain Measure (PPPM)^{10,11} at 12 and 24 hours after discharge. The PPPM is a 15-item instrument validated for children aged 1 to 12 years that examines the relationship between parent reports of child behaviors and child-rated pain in the postoperative period after minor surgery. Caregivers were contacted by telephone 24 hours after discharge, and families were asked to return the study journals by mail.

Sample Size and Statistical Analysis

A sample size of 25 in each treatment group provided 80% power to detect an effect size of 0.80 in the difference between FACES scores using the Mann-Whitney test with a Bonferroni-corrected 2-tailed significance level of .01^{12,13} (nQuery Advisor, version 7.0; Statistical Solutions). The effect size that we were sensitive to with the sample sizes was a difference in median FACES scores of 1 point or more between the 2 groups.

We used the Mann-Whitney test to compare FACES scores at the first obtainable score and at 10, 20, 30, and 40 or more minutes in the PACU and the PPPM scores at home. We compared the proportion of patients who had FACES pain scores of 0 between the groups using the 2-tailed Fisher exact test. Subsequent FACES scores were obtained every 4 hours (4, 8, 12, 16, 20, and 24) at home. Multivariable logistic regression was used to establish independent predictors of pain with age, body mass index (calculated as weight in kilograms divided by height in meters squared), and duration of surgery included as covariates.¹⁴ Vital signs (heart rate, mean arterial pressure, respiratory rate, and oxygen saturation level) were compared by the 2-tailed *t* test. Statistical analysis was performed using commercially available software (SPSS, version 19.0; SPSS Inc/IBM).

Results

Characteristics of Study Population

Among the 133 patients undergoing umbilical herniorrhaphy at Boston Children's Hospital from November 2009 through May 2011, 100 were deemed eligible for inclusion in the study (Figure 1). Thirty-three patients did not meet inclusion criteria owing to the need for additional concomitant surgery (n = 14), a coexisting medical condition categorized as American Society of Anesthesiologists classification of III or IV (n = 4), a history of a bleeding disorder (n = 1), being a ward of the state (n = 6), being non-English or non-Spanish speaking (n = 3), receiving preoperative analgesic within 24 hours before sur-

Table 1. Demographic and Clinical Characteristics of 52 Patients With Umbilical Hernia Enrolled in the Study^a

Characteristic	LAI (n = 25)	BRSB (n = 27)	P Value ^b
Age, mean (SD), y	6.1 (2.4)	6.0 (2.5)	.95
Male sex, %	14 (56)	12 (44)	.41
Weight, mean (SD), kg	22.3 (8.0)	26.3 (13.3)	.20
BMI, mean (SD)	16.1 (1.8)	18.8 (3.9)	.02
ASA class, No. (%)			
I	18 (72)	21 (78)	.63
II	7 (28)	6 (22)	
Race, No. (%)			
White	13 (52)	7 (26)	.22
Black	6 (24)	7 (26)	
Other	1 (4)	2 (7)	
Unknown	5 (20)	11 (41)	
Insurance, No. (%)			
Private	21 (84)	21 (78)	.57
Public	4 (16)	6 (22)	
Length of surgery, mean (SD), min	30 (9)	39 (18)	.04
Time from PACU arrival to first obtainable pain score, mean (SD), min	33.9 (20.8)	36.5 (24.1)	.68
Length of PACU stay, mean (SD), min	88 (34)	81 (38)	.38

Abbreviations: ASA, American Society of Anesthesiologists; BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); BRSB, bilateral rectus sheath block; LAI, local anesthetic infiltration; PACU, postanesthesia care unit.

^a All patients received ropivacaine hydrochloride.

^b Statistically significant P values are given in boldface type. The χ^2 test was used for categorical data and the 2-tailed t test for means.

gery (n = 3), and inability to express pain using FACES (n = 2). Forty-two families refused to participate and 6 were not successfully approached. Fifty-two patients were successfully randomized, and informed consent was obtained (Figure 1).

Characteristics of Study Participants

Patients were matched for age and sex and randomized to receive LAI (n = 25) or BRSB (n = 27). Table 1 shows no significant group differences in age, sex, weight, American Society of Anesthesiologists classification, race, insurance, time from PACU arrival to the first obtainable FACES score, or length of PACU stay. The body mass index was significantly higher (18.8 vs 16.1; $P = .02$) and surgical case duration was significantly longer (39 vs 30 minutes; $P = .04$) in the BRSB group compared with the LAI group. Preoperative vital signs (heart rate, mean arterial pressure, respiratory rate, and oxygen saturation level) and vital signs obtained on arrival in the PACU were not significantly different between the 2 groups (data not shown). No adverse events requiring immediate medical attention associated with the surgical procedure or the postoperative course were reported in either group.

FACES Scores Reported in the PACU

Table 2 shows that the preoperative FACES scores for all study participants were 0. Median FACES scores were significantly lower in the BRSB group at 10 ($P = .04$), 30 ($P = .01$), and 40 minutes or longer ($P = .03$) after arrival in the PACU compared with the LAI group. By classifying FACES scores into groups of 0 (no pain) and 1 to 5 (pain), treatment type was confirmed as an independent significant predictor of pain in the PACU at 10, 30, and 40 minutes or longer after arrival independent of age, body mass index, or duration of surgery. The percentage of children reporting no pain (FACES score, 0) was significantly higher in the BRSB group compared with the LAI group at these 3 times

(Figure 2). A total of 9 of the 23 patients in the LAI group (39%) and 16 of 24 in the BRSB group (67%) had no pain (FACES score of 0) at 10 minutes in the PACU; this difference of 28% has a 95% CI of 5% to 50%. At 30 minutes, 6 of 20 patients in the LAI group (30%) vs 15 of 21 in the BRSB group (71%) had no pain (41% difference [95% CI, 15%-62%]). At 40 minutes or longer, 10 of 19 patients in the LAI group (53%) and 9 of 10 in the BRSB group (90%) had no pain (38% difference [95% CI, 12%-57%]).

FACES and PPPM Scores Reported at Home

We found no significant differences in FACES scores obtained at any of the 4-hour intervals within the first 24 hours after discharge between the 2 groups (Table 2). Table 2 also shows no significant group differences in PPPM scores obtained at 12 and 24 hours after discharge.

Opioid and Nonopioid Medications

We found no significant differences in use of opioid and non-opioid medications between the treatment groups at any time in the PACU or at home (data not shown). However, when examining total use of analgesic medications in the PACU, we found fewer doses of opioid and nonopioid (acetaminophen) medications given to the BRSB group (5 opioid and 5 non-opioid) compared with the LAI group (11 opioid and 10 non-opioid) (data not shown). This trend was not noted at home, where 65 total doses of analgesics (20 opioid and 45 non-opioid) were given to the LAI group, and 63 total doses of analgesics (24 opioid and 39 nonopioid) were given to the BRSB group. We examined compliance with the study protocol for analgesic use at home; of 63 analyzable data points, 30 (48%) were compliant, 26 (41%) received stronger analgesics than indicated, and 7 (11%) received weaker analgesics than indicated. All data points with a FACES score of 0 (n = 2) or 1 (n = 12) at home received stronger analgesics than indicated.

Table 2. Comparison of Pain Scores Between the Treatment Groups

Measure, Defined Interval	LAI Group		BRSB Group		P Value ^a
	Score, Median (Range)	No. of Patients	Score, Median (Range)	No. of Patients	
Preoperative FACES	0 (0-0)	25	0 (0-0)	27	>.99
PACU, FACES					
First obtainable	0 (0-5)	25	0 (0-4)	27	.35
10 min	1 (0-5)	23	0 (0-5)	24	.04
20 min	1 (0-4)	18	0 (0-5)	26	.34
30 min	1 (0-5)	20	0 (0-5)	21	.01
≥40 min	1 (0-4)	19	0 (0-1)	10	.03
Home, FACES, h					
4	2 (0-5)	22	2 (0-5)	25	.36
8	2 (0-5)	18	2 (0-5)	20	.98
12	1 (0-4)	12	2 (0-5)	12	.27
16	1 (0-3)	12	1 (0-4)	10	.94
20	1 (0-5)	21	1 (0-5)	23	.56
24	1 (0-3)	21	1 (0-4)	24	.19
Home, PPPM, h					
12	7 (2-13)	21	9 (0-13)	25	.54
24	5 (0-12)	21	5 (0-11)	25	.96

Abbreviations: BRSB, bilateral rectus sheath block; FACES, FACES Pain Rating Scale; LAI, local anesthetic infiltration; PACU, postanesthesia care unit; PPPM, Parent's Postoperative Pain Measure.

^a Statistically significant P values are given in boldface (calculated by nonparametric Mann-Whitney test).


Discussion

Patients' perceptions of their hospital care have been extensively studied, especially in adults, with an important finding that improvements are needed in pain management.¹⁵ These studies have led to the creation of acute pain management services in many tertiary hospitals with the aim of increasing patient satisfaction by providing adequate pain relief, cutting costs by reducing the length of hospitalization due to adverse side effects associated with increased opioid analgesic use, and decreasing the incidence of chronic pain.¹⁶ Similar data on parents' perceptions of postoperative pain control in children are not well known.

Use of regional anesthesia techniques for abdominal operations in children is being increasingly used with sparse data about efficacy. Previous studies since the 1980s have examined regional anesthesia for pediatric inguinal and scrotal procedures with variable postoperative analgesia after ilioinguinal nerve block.^{17,18} Our study differs from previous trials examining related questions in similar patient populations in that 1 prior pilot study by Isaac et al³ did not include ultrasonographic guidance to perform BRSB. A more recent study by Gurnaney et al⁸ examined analgesic use between the treatment groups as a primary objective; the BRSB group received the block before the incision, whereas the LAI group received a local anesthetic after wound closure. We compared FACES pain scores at defined intervals in the PACU and at home within the first 24 hours after surgery.

Our study has shown that the BRSB group has a significantly higher percentage of patients with a FACES score of 0 (ie, no pain) and lower FACES scores at multiple time points in the PACU compared with the LAI group. We also found that the total use of analgesic (opioid and nonopioid) medications in the PACU was reduced in the BRSB group compared with

Figure 2. Percentage of Patients With No Pain (Indicates FACES Pain Rating Scale Score of 0) in the Postanesthesia Care Unit (PACU)



*P < .05, 2-tailed Fisher exact test. BRSB indicates bilateral rectus sheath block; LAI, local anesthetic infiltration.

the LAI group. At each time point in the PACU and at home, we did not find a significant difference in opioid or nonopioid medication use. The clinical significance of these findings is encouraging with regard to efficacy of ultrasonography-guided BRSB in providing short-term postoperative analgesia after umbilical hernia repair. We presume that 1 benefit of the BRSB from the patient's and the family's standpoints is that no pain (FACES score of 0) is more desirable than any level of pain (any FACES score >0). Differences in FACES scores between the study groups may have been minimized by the routine use of intraoperative ketorolac for all patients at our institution, which was mandated by the surgeons. Lack of prolonged analgesic effects of the BRSB at home is unfortunate, but these findings may be owing partly to additional

methodologic challenges. Unsworth et al¹⁹ used a similar self-reported pain scale strategy for administration of postoperative analgesics at home in children undergoing adenotonsillectomy. When FACES scores were obtained at defined times in the first 3 days after surgery, a similar amount of analgesic was administered to the children when compared with parent administration of analgesics without the use of FACES scores. They noted that analgesic administration was concordant with children's pain scores in 68% of administrations, which is higher than the 48% noted in our study. These findings underscore the complicated dynamics involved in parental perception of their child's postoperative pain and their subsequent attempts to address that pain. Compliance is altered if some parents, regardless of pain scores, prefer to administer stronger opioids or forgo opioid use for fear of possible adverse effects. We observed both scenarios in our study.

Our study findings may be generalizable to patients undergoing single-incision laparoscopic surgery (SILS) for abdominal operations using an umbilical port. A recent prospective, randomized trial by St Peter et al²⁰ noted that the SILS group received a greater total dose of analgesics during their hospital stay compared with the 3-port group. Two meta-analyses comparing conventional laparoscopic cholecystectomy with SILS cholecystectomy suggest that considerable postoperative pain remains after SILS cholecystectomy.^{21,22} Future studies examining the role of ultrasonography-guided BRSB in reducing postoperative pain after adult and pediatric SILS abdominal surgery may provide additional evidence.

Our study has some limitations. First, although we used a randomized clinical trial design, the treatment groups are not large and power is thus limited in assessing differences in opioid use between the groups. Second, postoperative pain after umbilical hernia repair may be widely variable, making detection of small changes in FACES scores difficult to capture. Third, our compliance data with administration of postoperative analgesia at home by parents based on the study protocol suggest that more work is needed to understand the complexities of parental perception of their child's postoperative pain, some of which may be mitigated by improved parental education.

Future studies could examine the use of longer-acting local anesthetic agents for use with ultrasonography-guided BRSB. We chose ropivacaine, which has been shown to have superior postoperative analgesic effects compared with lidocaine when administered as a pediatric ilioinguinal nerve block.²³ Longer-acting local anesthetic agents still in development, such as neosaxitoxin,^{24,25} may demonstrate improved postoperative analgesia at home and in the PACU. The addition of clonidine hydrochloride to the ropivacaine blocks may further extend the analgesic effects.²⁶ In addition, administering the anesthetic before surgical incision for both treatment arms would allow one to study the role of preemptive analgesia in reducing postoperative pain in the PACU after each of the treatment arms. As newer pediatric regional anesthetic techniques are developed, studies should be performed to examine the efficacy of ultrasonography-guided BRSB with ultrasonography-guided bilateral transversus abdominis plane blocks²⁷ in reducing postoperative pain after open and laparoscopic pediatric abdominal surgical procedures requiring midline and lateral incisions.

Finally, as with all new technologies, whether the cost of the BRSB justifies the clinical benefit it provides needs further study. The current methods used for this study, however, are not suitable for this analysis. First, we originally hoped to administer the anesthetic before surgical incision for both treatment arms to study the role of preemptive analgesia in reducing postoperative pain in the PACU. However, because surgical preference required the local anesthesia to be delivered at the conclusion of surgery, we designed the BRSB arm to mirror this, thereby inflating the total length of surgery for the BRSB arm because the block could only be performed after the surgical dressings were applied. Second, patients were not billed for the BRSB in this study, and we therefore do not have any accurate cost data for that arm. Third, a future study comparing BRSB with LAI in providing preemptive analgesia could compare postoperative opioid use and frequency of opioid-induced adverse effects that may prolong recovery. Decreased PACU stay or avoidance of hospital admission could reduce costs.

ARTICLE INFORMATION

Accepted for Publication: November 6, 2012.

Published Online: June 12, 2013.

doi:10.1001/jamasurg.2013.1442.

Author Affiliations: Department of Anesthesiology, Perioperative and Pain Medicine, Boston Children's Hospital, Harvard Medical School, Boston, Massachusetts (Dingeman, Chung, Clendenin, Lee); Department of Surgery, Boston Children's Hospital, Harvard Medical School, Boston, Massachusetts (Barus, Tracy, Dennett, Zurakowski, Chen); Division of Biostatistics, Boston Children's Hospital, Harvard Medical School, Boston, Massachusetts (Johnson, Zurakowski); now with the Department of Anesthesia, Cohen Children's Medical Center of New York, North Shore-Long Island Jewish Health System, New Hyde Park, New York (Dingeman); now with the Department of Anesthesia, University of Massachusetts Memorial Medical Center, Worcester, Massachusetts (Chung); now with the

Division of Anesthesia, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio (Lee); now with Meridian Health, Neptune City, New Jersey (Johnson); now a medical student at University of Massachusetts Medical School, Worcester, Massachusetts (Tracy).

Author Contributions: *Study concept and design:* Dingeman, Tracy, Zurakowski, and Chen.

Acquisition of data: Dingeman, Barus, Chung, Clendenin, Lee, Tracy, Dennett, and Chen.

Analysis and interpretation of data: Dingeman, Barus, Johnson, Dennett, Zurakowski, and Chen.

Drafting of the manuscript: Dingeman, Barus, Johnson, Dennett, Zurakowski, and Chen.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Zurakowski.

Obtained funding: Dingeman, Tracy, and Chen.

Administrative, technical, and material support: Dingeman and Chen.

Study supervision: Dingeman, Barus, Zurakowski, and Chen.

Conflict of Interest Disclosures: None reported.

Funding/Support: This study was supported by a pilot grant from Harvard Catalyst/The Harvard Clinical and Translational Science Center, Boston Children's Hospital Surgical Foundation, and the Department of Anesthesiology, Perioperative, and Pain Medicine.

Previous Presentation: This study was presented at the 65th Annual Postgraduate Assembly in Anesthesiology; December 10, 2011; New York, NY.

Additional Contributions: Terry L. Buchmiller, MD, Steven J. Fishman, MD, Tom Jaksic, MD, PhD, Craig W. Lillehei, MD, Bradley C. Linden, MD, Konstantinos Papadakis, MD, Mark Puder, MD, PhD, Shawn J. Rangel, MD, MSCE, Robert C. Shamberger, MD, C. Jason Smithers, MD, Christopher B. Weldon, MD, PhD, and Jay M. Wilson, MD, enrolled patients. Samantha Butler, PhD, participated on the Data

Safety Monitoring Board. Clarissa Valim, MD, ScD, Susan McDermott, MPH, RN, Lin Huang, PhD, and Adam Simmons, MPH, assisted with randomization. Kelly Hermance-Perry, RN, and Rebecca Brennan, MS, helped with scheduling.

REFERENCES

- Ferguson S, Thomas V, Lewis I. The rectus sheath block in paediatric anaesthesia: new indications for an old technique? *Paediatr Anaesth*. 1996;6(6):463-466.
- Courreges P, Poddevin F, Lecoutre D. Para-umbilical block: a new concept for regional anaesthesia in children. *Paediatr Anaesth*. 1997;7(3):211-214.
- Isaac LA, McEwen J, Hayes JA, Crawford MW. A pilot study of the rectus sheath block for pain control after umbilical hernia repair. *Paediatr Anaesth*. 2006;16(4):406-409.
- Willschke H, Marhofer P, Bösenberg A, et al. Ultrasonography for ilioinguinal/iliohypogastric nerve blocks in children. *Br J Anaesth*. 2005;95(2):226-230.
- Willschke H, Bösenberg A, Marhofer P, et al. Ultrasonographic-guided ilioinguinal/iliohypogastric nerve block in pediatric anesthesia: what is the optimal volume? *Anesth Analg*. 2006;102(6):1680-1684.
- de Jose Maria B, Götzens V, Mabrok M. Ultrasound-guided umbilical nerve block in children: a brief description of a new approach. *Paediatr Anaesth*. 2007;17(1):44-50.
- Willschke H, Bösenberg A, Marhofer P, et al. Ultrasonography-guided rectus sheath block in paediatric anaesthesia: a new approach to an old technique. *Br J Anaesth*. 2006;97(2):244-249.
- Gurnaney HG, Maxwell LG, Kraemer FW, Goebel T, Nance ML, Ganesh A. Prospective randomized observer-blinded study comparing the analgesic efficacy of ultrasound-guided rectus sheath block and local anaesthetic infiltration for umbilical hernia repair. *Br J Anaesth*. 2011;107(5):790-795.
- Wong DL, Baker CM. Pain in children: comparison of assessment scales. *Pediatr Nurs*. 1988;14(1):9-17.
- Chambers CT, Reid GJ, McGrath PJ, Finley GA. Development and preliminary validation of a postoperative pain measure for parents. *Pain*. 1996;68(2-3):307-313.
- Chambers CT, Finley GA, McGrath PJ, Walsh TM. The parents' postoperative pain measure: replication and extension to 2-6-year-old children. *Pain*. 2003;105(3):437-443.
- Noether GE. Sample size determination for some common nonparametric tests. *J Am Stat Assoc*. 1987;82(398):645-647.
- Browner WS, Newman TB, Cummings SR, Hulley SB. Estimating sample size and power: the nitty-gritty. In: Hulley SB, Cummings SR, Browner WS, Grady D, Hearst N, Newman TB, eds. *Designing Clinical Research*. 2nd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2001:65-91.
- Katz MH. *Multivariable Analysis: A Practical Guide for Clinicians and Public Health Researchers*. 3rd ed. New York, NY: Cambridge University Press; 2011:74-87.
- Jha AK, Orav EJ, Zheng J, Epstein AM. Patients' perception of hospital care in the United States. *N Engl J Med*. 2008;359(18):1921-1931.
- Lee A, Chan SK, Chen PP, Gin T, Lau AS, Chiu CH. The costs and benefits of extending the role of the acute pain service on clinical outcomes after major elective surgery. *Anesth Analg*. 2010;111(4):1042-1050.
- Shandling B, Steward DJ. Regional analgesia for postoperative pain in pediatric outpatient surgery. *J Pediatr Surg*. 1980;15(4):477-480.
- Kokinsky E, Thornberg E, Östlund AL, Larsson LE. Postoperative comfort in paediatric outpatient surgery. *Paediatr Anaesth*. 1999;9(3):243-251.
- Unsworth V, Franck LS, Choonara I. Parental assessment and management of children's postoperative pain: a randomized clinical trial. *J Child Health Care*. 2007;11(3):186-194.
- St Peter SD, Adibe OO, Juang D, et al. Single incision versus standard 3-port laparoscopic appendectomy: a prospective randomized trial. *Ann Surg*. 2011;254(4):586-590.
- Garg P, Thakur JD, Garg M, Menon GR. Single-incision laparoscopic cholecystectomy vs conventional laparoscopic cholecystectomy: a meta-analysis of randomized controlled trials. *J Gastrointest Surg*. 2012;16(8):1618-1628.
- Markar SR, Karthikesalingam A, Thrumurthy S, Muirhead L, Kinross J, Paraskeva P. Single-incision laparoscopic surgery (SILS) vs conventional multiport cholecystectomy: systematic review and meta-analysis. *Surg Endosc*. 2012;26(5):1205-1213.
- Tsuchiya N, Ichizawa M, Yoshikawa Y, Shinomura T. Comparison of ropivacaine with bupivacaine and lidocaine for ilioinguinal block after ambulatory inguinal hernia repair in children. *Paediatr Anaesth*. 2004;14(6):468-470.
- Rodríguez-Navarro AJ, Berde CB, Wiedmaier G, et al. Comparison of neosaxitoxin versus bupivacaine via port infiltration for postoperative analgesia following laparoscopic cholecystectomy: a randomized, double-blind trial. *Reg Anesth Pain Med*. 2011;36(2):103-109.
- Wylie MC, Johnson VM, Carpino E, et al. Respiratory, neuromuscular, and cardiovascular effects of neosaxitoxin in isoflurane-anesthetized sheep. *Reg Anesth Pain Med*. 2012;37(2):152-158.
- Cucchiari G, Ganesh A. The effects of clonidine on postoperative analgesia after peripheral nerve blockade in children. *Anesth Analg*. 2007;104(3):532-537.
- Suresh S, Chan VW. Ultrasound guided transversus abdominis plane block in infants, children and adolescents: a simple procedural guidance for their performance. *Paediatr Anaesth*. 2009;19(4):296-299.

Invited Commentary

Pain Control After Umbilical Hernia Repair How Difficult Can We Make It?

Brad W. Warner, MD

Important to the overall experience for a child undergoing elective surgery is effective pain management. In this excellent study by Dingeman et al,¹ a randomized prospective trial compared infiltration of local anesthetic with an ultrasonography-guided bilateral rectus sheath block (BRSB) after umbilical hernia repair. The authors report that the overall pain scores were better with fewer supplemental pain medications needed in the BRSB group.

This report has some limitations. Given the frequency with which umbilical hernia repair is performed, this study would be more convincing if a larger number of patients were enrolled. Further, although statistically significant, the pain score

differences between the 2 groups were generally on the order of a single point. Therefore, whether these differences were biologically important remains unclear. Because preemptive analgesia is recognized to be more effective in minimizing postoperative pain, an even greater effect of the BRSB is likely if administered before the incision. Thus, a better study design would compare preincision BRSB with the traditional postrepair infiltration of local anesthetic.

The cost difference between local anesthesia and BRSB was not mentioned and warrants justification, especially in face of the modest difference in pain scores between groups. Injecting local anesthetic is easy and does not require specific equipment. The BRSB procedure requires an



Related article page 707

ultrasonographic machine and specific training to perform this technique successfully. Cost may therefore be higher as a result of the purchase and maintenance of the machine. An anesthesiologist charge for providing a regional anesthetic may be added. Further, the technique and training

required for BRSB will undoubtedly add to operative time. Finally, risks unique to BRSB need to be monitored. All these factors must be considered because they undoubtedly add complexity to a straightforward and typically well-tolerated procedure.

ARTICLE INFORMATION

Author Affiliation: Division of Pediatric Surgery, Department of Surgery, Washington University School of Medicine, St Louis Children's Hospital, St Louis, Missouri.

Corresponding Author: Brad W. Warner, MD, Division of Pediatric Surgery, Department of

Surgery, Washington University School of Medicine, St Louis Children's Hospital, Children's Place, Ste 5540, St Louis, MO 63110 (brad.warner@wustl.edu)

Published Online: June 12, 2013.
doi:10.1001/jamasurg.2013.1453.

Conflict of Interest Disclosures: None reported.

REFERENCE

1. Dingeman RS, Barus LM, Chung HK, et al. Ultrasonography-guided bilateral rectus sheath block vs local anesthetic infiltration after pediatric umbilical hernia repair: a prospective randomized clinical trial [published online June 12, 2013]. *JAMA Surg*. 2013;148(8):707-713.