

# Establishing Standards of Quality for Elderly Patients Undergoing Pancreatic Resection

Wande B. Pratt, MD, MPH; Anupama Gangavati, MD; Kathryn Agarwal, MD; Robert Schreiber, MD; Lewis A. Lipsitz, MD; Mark P. Callery, MD; Charles M. Vollmer Jr, MD

**Objective:** To evaluate pancreatic surgery as a model for high-acuity surgery in elderly patients for immediate and long-term outcomes, predictors of adverse outcomes, and hospital costs.

**Design:** Retrospective case series.

**Setting:** University tertiary care referral center.

**Patients:** Four hundred twelve consecutive patients who underwent pancreatic resection from October 1, 2001, through March 31, 2008, for benign and malignant periampullary conditions.

**Main Outcome Measures:** Clinical outcomes were compared for elderly ( $\geq 75$  years) and nonelderly patient cohorts. Quality assessment analyses were performed to show the differential impact of complications and resource utilization between the groups.

**Results:** The elderly cohort constituted one-fifth of all

patients. Benchmark standards of quality were achieved in this group, including low operative mortality (1%). Despite higher patient acuity, clinical outcomes were comparable to those of nonelderly patients at a marginal cost increase (median, \$2202 per case). Cost modeling analysis showed further that minor and moderate complications were more frequent but no more debilitating for elderly patients. Major complications, however, were far more threatening to older patients. In these cases, duration of hospital stay doubled, and invasive interventions were more commonly deployed.

**Conclusions:** Quality standards for pancreatic resection in the elderly can—and should—mirror those for younger patients. Age-related care, including geriatric consultation, supplemental enteral nutrition, and early rehabilitation placement planning, can be designed to mitigate the impact of complications in the elderly and guarantee quality.

*Arch Surg.* 2009;144(10):950-956

**T**HE ELDERLY CONSTITUTE THE fastest-growing age demographic in the United States. The US Census Bureau estimates that 12% of the population is currently 65 years or older. By 2030, this demographic will double in absolute number and represent 20% of the US population.<sup>1</sup> Longer life expectancy, coupled with advancements in surgical techniques, has resulted in more elderly patients presenting for surgery.<sup>2</sup> Surgeons and other physicians will wrestle with reasonable indications for surgery and its effect on patient morbidity, quality of life, resource utilization, and expenditures for this cohort.

These dilemmas are common when patients present with periampullary malignant neoplasms. Pancreatic cancer is among the deadliest cancers, with annual mortality approximating its incidence. Beyond mortality, the burden of this aggressive cancer is overwhelming—often presenting late in its natural course and lacking any effective screening test.<sup>3-6</sup> Surgical resection remains the only po-

tential curative treatment but is an option for only 10% to 25% of patients.<sup>7</sup> Although some randomized trials have shown a marginal survival benefit when adjuvant therapy is used,<sup>8</sup> chemoradiotherapy requires prompt and complete surgical recovery. This option is precluded in some patients, perhaps more so in the elderly, owing to delayed recovery from postoperative complications, low physiologic reserve, or early mortality.<sup>9</sup>

Standards for surgical care are receiving increasing scrutiny, particularly in high-acuity procedures such as pancreatic resection, in which patients are typically older, baseline patient risk is high, and complications are frequent, costly, and debilitating. New process and systems improvement initiatives rely on established quality indicators, the assessment of practice outcomes, and standardized perioperative management approaches.<sup>10-18</sup> However, it remains unclear whether contemporary standards of quality for pancreatic resection are equivalent for young and elderly patients and to what extent such approaches can optimize outcomes

**Author Affiliations:** Department of Surgery (Drs Pratt, Callery, and Vollmer), and Division of Gerontology, Department of Medicine (Drs Gangavati, Agarwal, and Lipsitz), Harvard Medical School, and Hebrew SeniorLife, Institute for Aging Research (Drs Schreiber and Lipsitz), Beth Israel Deaconess Medical Center, Boston, Massachusetts.

for elderly patients. This analysis addresses these uncertainties and provides recommendations for achieving optimal outcomes when pancreatic resection is performed in older patients.

## METHODS

### PATIENTS, OPERATIONS, AND MANAGEMENT

Two fellowship-trained pancreatobiliary surgical specialists (M.P.C. and C.M.V.) performed 412 consecutive pancreatic resections from October 1, 2001, through March 31, 2008, including 272 proximal, 124 distal, 9 central, and 7 total pancreatectomies, for a full spectrum of benign and malignant periampullary pathological findings. For pancreatoduodenectomy, the most common indication was pancreatic ductal adenocarcinoma (109 patients [40%]), whereas distal pancreatectomy was performed most often for cystic neoplasms (54 [44%]). Central pancreatectomy was selectively applied to lesions of the body and neck for its ability to preserve pancreatic endocrine and exocrine function. Total pancreatectomy was reserved for lesions not amenable to limited pancreatic resection, most often for intraductal papillary mucinous neoplasms.

The Carepath for Pancreatic Resection, a standardized clinical pathway for perioperative care, was used at our institution. This pathway, previously described in detail elsewhere, outlines a reproducible multidisciplinary approach to preoperative, intraoperative, and postoperative management.<sup>17,18</sup> Geriatric consultation and specific age-related care were typically provided for patients older than 75 years. This includes evaluation of comorbid conditions, particularly cardiac and respiratory illness. Gerontologists also identify family support structures and discuss end-of-life and resuscitation preferences. Nutritional evaluation and cognitive and functional analyses are regularly performed. In the postoperative period, geriatric consultation is useful for fluid management, pain control, and prevention and treatment of postoperative delirium. These specific interventions were formally incorporated into our care pathway in 2006 but were frequently used in earlier years.

### DATA COLLECTION

Data were prospectively collected in accordance with guidelines for human subjects research, as approved by our institutional review board. Recorded variables have been described and defined elsewhere,<sup>17-19</sup> including analysis of preoperative status with the American Society of Anesthesiologists (ASA) Classification of Physical Status<sup>20</sup> and the Karnofsky scale.<sup>21</sup> Patients were classified as elderly (aged  $\geq 75$  years) or nonelderly. Many gerontologists and pancreatic surgeons now regard patients older than 75 years as elderly, particularly because the average age for patients undergoing pancreatic resection is 63 years.<sup>14</sup>

### TRADITIONAL QUALITY ASSESSMENTS

Clinical and economic outcomes were compared between cohorts. In addition to accepted baseline indicators of quality—complications, duration of hospital stay, and perioperative mortality—we explored other emerging measures of surgical quality, which have been defined previously.<sup>17</sup> The incidence and severity of complications were defined according to the Clavien scheme, which describes 5 grades of clinical severity, based on escalating levels of therapeutic interventions required.<sup>22</sup> Briefly, grades 1 and 2 events represent complications of minor sever-

ity, and grade 3A reflects a moderate complication. Grades 3B and 4 represent major complications, and grade 5 represents death.

## CONTEMPORARY QUALITY ASSESSMENTS

Beyond these traditional assessments, 2 other analyses—ratio of observed to expected (O:E) morbidity and deviation-based cost modeling (DBCM)—were performed to measure the incremental impact of increasing age on outcomes and to determine whether elderly status significantly alters the anticipated postoperative clinical course. The rationale and details of these analyses have been published previously.<sup>17-19,23</sup> Using the actual incidence of postoperative complications and the mean Physiologic and Operative Severity Score for the Enumeration of Mortality and Morbidity (POSSUM),<sup>19,23</sup> we evaluated surgical quality through variance in O:E morbidity. A ratio equal to 1.00 demonstrates expected performance. Ratios of greater than 1.00 suggest that the outcomes are worse than expected. Conversely, ratios of less than 1.00 suggest that the outcomes achieved are better than expected. Deviations from our standard care pathway were evaluated using DBCM to compare the clinical and economic impact of complications.<sup>18</sup> For any given operation or procedure, DBCM links the severity of complications with the length of stay. Hospital costs for each deviation class were compared using a summary measure of cost efficiency that combines the relative proportion of each deviation category (on course to major) with its median hospital cost.<sup>18</sup> This provides the weighted average median cost per patient, reflecting the direct contribution of complications to costs.

### STATISTICAL ANALYSIS

Categorical variables were compared using the  $\chi^2$ , Fisher exact, and univariate logistic regression statistics; continuous variables were compared using analysis of variance, unpaired *t* tests for independent variables, and simple linear regression. Differences between O:E morbidity were assessed using the Hosmer-Lemeshow  $\chi^2$  goodness-of-fit method.<sup>24</sup> Factors associated with morbidity were calculated by cross-tabulation using  $\chi^2$  and Pearson correlation tests. We considered  $P < .05$  to be statistically significant. Computations were performed using commercially available software (SPSS 14.0 [SPSS, Inc, Chicago, Illinois] and Stata 8.2 [StataCorp LP, College Station, Texas]).

## RESULTS

### BASELINE CHARACTERISTICS

From October 1, 2001, through March 31, 2008, pancreatic resection was performed in 76 elderly patients (18%) and 336 nonelderly patients (82%). Elderly patients had significantly lower preoperative albumin levels and median body mass indexes (**Table 1**). Several measures of patient acuity were significantly worse among the elderly patients. Elderly patients were twice as likely to present with American Society of Anesthesiologists classifications of III or IV (odds ratio, 2.17; 95% confidence interval [CI], 1.28-3.68;  $P = .004$ ), and the median POSSUM physiologic severity score was significantly higher (25 vs 17;  $P < .001$ ). Karnofsky scores of 80 or 70 were more frequent, but resections were not performed for patients with scores of less than 70. Comorbidity rates, except for chronic obstructive pulmonary disorder, were equivalent. Elderly patients harbored malignant neoplasms significantly more of-

**Table 1. Preoperative Patient Characteristics for Nonelderly and Elderly Patients<sup>a</sup>**

Preoperative Characteristics	Nonelderly Cohort (n=336)	Elderly Cohort (n=76)	P Value
Age, median (range), y	58 (23-74)	79 (75-90)	<.001
Sex			
Male	161 (48)	33 (43)	.28
Female	175 (52)	43 (57)	
BMI, median (range)	26.0 (15.4-48.9)	24.4 (17.5-37.8)	.005
Albumin level, median (range), g/dL	3.9 (1.4-5.1)	3.6 (1.4-4.6)	.002
Karnofsky score <sup>b</sup>			
90 (Able to perform normal activity with only minor symptoms)	256 (76)	38 (50)	<.001
80 (Able to perform normal activity with effort, some symptoms)	70 (21)	31 (41)	
70 (Able to care for self, but unable to do normal activities)	10 (3)	7 (9)	
ASA physical status			
Class I	5 (1)	0	.02
Class II	163 (49)	24 (32)	
Class III	161 (48)	48 (63)	
Class IV	7 (2)	4 (5)	
POSSUM physiologic severity score (range) <sup>c</sup>	17 (12-39)	25 (15-40)	<.001
Coronary heart disease	44 (13)	19 (25)	.56
Hypertension	145 (43)	42 (55)	.053
COPD	21 (6)	13 (17)	.004
Diabetes mellitus	69 (21)	20 (26)	.17
Final pathological finding			
Benign	168 (50)	12 (16)	<.001
Malignant	168 (50)	64 (84)	

Abbreviations: ASA, American Society of Anesthesiologists; BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); COPD, chronic obstructive pulmonary disease; POSSUM, Physiologic and Operative Severity Score for the Enumeration of Mortality and Morbidity.

SI conversion factor: To convert albumin to grams per liter, multiply by 10.

<sup>a</sup>Unless otherwise indicated, data are expressed as number (percentage) of patients.

<sup>b</sup>Karnofsky Performance Status.<sup>21</sup>

<sup>c</sup>Described in Copeland et al.<sup>23</sup>

**Table 2. Operative Outcomes for Nonelderly and Elderly Patients After Pancreatic Resection<sup>a</sup>**

Intraoperative Outcomes	Nonelderly Patients (n=336)	Elderly Patients (n=76)	P Value
Resection types			
Pancreatoduodenectomy	213 (63)	59 (78)	.09
Distal pancreatectomy	108 (32)	16 (21)	
Central pancreatectomy	9 (3)	0	
Total pancreatectomy	6 (2)	1 (1)	
Operative time, median (range), min	358 (76-780)	361 (133-628)	.73
Blood loss, median (range), mL	350 (0-15 000)	350 (50-1500)	.42
Blood transfusion	44 (13)	17 (22)	.03
POSSUM operative severity score, median (range) <sup>b</sup>	15 (13-27)	15 (13-23)	.55

Abbreviation: POSSUM, Physiologic and Operative Severity Score for the Enumeration of Mortality and Morbidity.

<sup>a</sup>Unless otherwise indicated, data are expressed as number (percentage) of patients.

<sup>b</sup>Described in Copeland et al.<sup>23</sup>

ten (64 of 76 patients [84%] vs 168 of 336 [50%];  $P < .001$ ), with pancreatic adenocarcinoma being the most common indication for nonelderly (96 patients [29%]) and elderly (35 [46%]) patients. Nonelderly patients more often presented with pancreatitis or other benign conditions (12 patients [16%] vs 168 [50%]).

## OPERATIVE FACTORS

Operative outcomes were statistically equivalent between the age groups (**Table 2**). Blood loss (median, 350 mL) and operative time were benchmarks for both cohorts, whereas the number of individuals receiving blood

transfusions was significantly greater among elderly patients. Surgical performance (POSSUM operative severity score) was equal, indicating that, despite higher patient acuity among elderly patients, equivalent operative conduct can be achieved.

## TRADITIONAL QUALITY ANALYSIS

### Elderly Patients

Among the 76 elderly patients, 55 (72%) developed at least 1 complication of any severity, but two-thirds of these (39 of 55) were minor in scope. Oliguria/hypotension (37

**Table 3. Postoperative Outcomes for Nonelderly and Elderly Patients After Pancreatic Resection<sup>a</sup>**

Postoperative Outcomes	Nonelderly Patients (n=336)	Elderly Patients (n=76)	P Value
Morbidity	161 (48)	55 (72)	<.001
Mortality	4 (1)	1 (1)	.64
Severity of complications			
None	175 (52)	21 (28)	.006
Grade 1	73 (22)	29 (38)	
Grade 2	39 (12)	10 (13)	
Grade 3	29 (9)	10 (13)	
Grade 4	16 (5)	5 (7)	
Grade 5	4 (1)	1 (1)	
Therapeutic interventions			
Antibiotics	88 (26)	30 (39)	.02
Parenteral nutrition	37 (11)	14 (18)	.03
Blood transfusion	44 (13)	26 (34)	<.001
Invasive interventions			
CT-guided percutaneous drainage	15 (4)	6 (8)	.17
Reoperation	12 (4)	8 (11)	.02
ICU utilization	18 (5)	3 (4)	.44
Duration of hospital stay, median, d	8	9	<.001
Discharge disposition <sup>b</sup>			
Home	218 (66)	19 (25)	<.001
Home with nursing assistance	95 (29)	22 (29)	
Rehabilitation facility	19 (6)	34 (45)	
Readmission	50 (15)	11 (14)	.556

Abbreviations: CT, computed tomography; ICU, intensive care unit.

<sup>a</sup>Unless otherwise indicated, data are expressed as number (percentage) of patients. Because of rounding, percentages may not total 100.

<sup>b</sup>Data are from patients eligible for discharge. Four in-hospital deaths occurred in the nonelderly cohort and 1 in the elderly cohort.

patients [49%]), delirium (22 [29%] vs 30 nonelderly patients [9%];  $P < .001$ ), wound infections (16 patients [21%]), and respiratory distress (12 [16%]) were common. The median hospital duration was 9 days. Although most returned home (19 [25%]) or required visiting nursing assistance (22 [29%]), 34 patients (45%) required further treatment at rehabilitation facilities. Hospital readmission and reoperation were infrequent (11 patients [14%] and 8 [11%], respectively); only 1 elderly patient (1.3%) died (of myocardial infarction). The median total hospital cost for these patients was \$19 852 (range, \$9873-\$164 756).

### Elderly vs Nonelderly Patients

Clinical outcomes were compared directly between elderly and nonelderly patients (**Table 3**). Overall, postoperative complications were significantly greater among elderly patients, although this is explained by differences in rates of minor complications (39 elderly patients [51%] vs 112 nonelderly patients [33%];  $P = .003$ ). Moderate and major complications (grades 3-5) were equivalent (16 elderly patients [21%] vs 49 nonelderly patients [15%];  $P = .11$ ). As a result, more antibiotic therapy, blood transfusions, and parenteral nutrition were required by the elderly. Moreover, elderly patients showed a trend toward increasing rates of image-guided drain-

**Table 4. Economic Outcomes for Nonelderly and Elderly Patients Undergoing Pancreatic Resection**

Hospital Costs	Median Cost per Cohort, \$		P Value
	Nonelderly Patients	Elderly Patients	
Radiology	339	683	.21
Laboratory	511	587	.01
Pharmacy	777	968	.02
Transfusion	134	401	.08
ICU	768	1016	.02
Operating room	4374	4626	.85
Room	6628	8271	.001
Total <sup>a</sup>	17 650	19 852	.001
Cost increase	...	2202	...

Abbreviations: Ellipses, not applicable; ICU, intensive care unit.

<sup>a</sup>Data represent the median total cost for each group, not the sum of the cost centers listed.

age and required reoperation 3 times more often (odds ratio, 3.18; 95% CI, 1.25-8.07;  $P = .01$ ). These factors contributed to a 1-day increase in median hospital stay and higher rates of rehabilitation placement.

Detailed economic analysis demonstrates that pancreatic resection was somewhat more costly among elderly patients, translating to \$2202 (or roughly 12% more) per patient (**Table 4**). This cost increase largely reflects increased laboratory evaluations, pharmacologic interventions, duration of intensive care unit stay, and the additive costs of an additional day of hospitalization.

### Risk Factors for Postoperative Morbidity

Multivariate logistic regression analysis of the overall cohort confirmed advanced age ( $P = .03$ ) and intraoperative blood loss ( $P = .004$ ) as predictive factors associated with postoperative morbidity. Advanced age was associated with a 2-fold increase in the likelihood of morbidity (odds ratio, 2.05; 95% CI, 1.07-3.93;  $P = .03$ ). This risk-adjusted analysis also suggests that elderly patients, on average, incur a 2-day increase in hospital stay ( $\beta = 2.35$ ; 95% CI, 1.03-3.67;  $P < .001$ ) and a \$9800 increase in total hospital costs ( $\beta = \$9774$ ; 95% CI, \$4117-\$15 431;  $P < .001$ ).

When elderly patients were scrutinized separately on univariate analysis, type of resection, lower preoperative hematocrit level, presence of a malignant neoplasm, and operative time were associated with any level of morbidity. However, these were not significant after multivariate analysis. Complications were broken into 2 strata, minor and moderate/major, to determine which factors predict complication severity. There were no identifiable risk factors for minor complications. However, intraoperative blood loss was the only significant factor ( $P = .03$ ) for moderate/major complications.

### CONTEMPORARY QUALITY ANALYSIS

For the complete time frame, the overall O:E morbidity ratio was 0.94, steadily improving from 1.24 in 2001 to 0.87 currently.<sup>17</sup> The O:E ratios were calculated sepa-

**Table 5. Deviation-Based Cost Modeling for Comparison of Nonelderly and Elderly Patients**

Deviation-Based Cost Modeling <sup>a</sup>	Nonelderly Patients (n=336)	Elderly Patients (n=76)	P Value
Deviation mix, No. (%) of patients			
On course	216 (64)	34 (45)	.01
Minor deviation	47 (14)	18 (24)	
Moderate deviation	44 (13)	12 (16)	
Major deviation	29 (9)	12 (16)	
On course			
Duration of hospital stay, d	7	8	.03
Total hospital costs, \$	16 828	18 385	.93
Minor deviation			
Duration of hospital stay, d	9	9	.82
Total hospital costs, \$	19 755	18 346	.37
Moderate deviation			
Duration of hospital stay, d	13	15	.06
Total hospital costs, \$	25 631	31 441	.07
Major deviation			
Duration of hospital stay, d	9	15	.20
Total hospital costs, \$	27 326	71 812	.01
Duration of hospital stay (weighted average), d	8.2	10.4	...
Duration of stay increase, d	...	2.2	...
Hospital costs (weighted average), \$	19 296	28 873	...
Cost increase, \$	...	9577	...

Abbreviation: Ellipses, not applicable.

<sup>a</sup> Described in full detail by Vanounou et al.<sup>18</sup>

rately for the elderly and nonelderly patient cohorts. Although elderly patients demonstrated higher expected and observed morbidity rates (both  $P < .001$ ), calculated O:E ratios were equivalent. This ratio declined more significantly during the 6-year time frame among younger patients (1.27 to 0.84), whereas elderly patients experienced more modest, but real, improvements (1.15 to 0.93).

When DBCM was applied, the incidence and severity of deviations was substantially greater among elderly patients (**Table 5**). Within the on-course category, elderly patients had longer hospital stays by 1 day ( $P = .03$ ); however, costs were similar. These metrics were equivalent for minor deviations. Moderate deviations were only marginally more severe and costly. Major deviations, however, were far more consequential to older patients. Hospital duration nearly doubled, and staggering cost discrepancies resulted.

Collectively, these differential outcomes are reflected in the weighted-average median hospital duration and cost, which depicts the full impact of complications. The overall impact of complications in elderly patients was a 2.2-day increase in duration of hospital stay and a \$9577 cost increase per patient, quite consistent with our risk-adjusted model described in the "Traditional Quality Analysis" subsection.

#### COMMENT

The recent growth of the elderly population affords physicians and surgeons unprecedented opportunities to understand—and perhaps mitigate—the impact of age on

surgical recovery.<sup>15</sup> This is particularly relevant when high-acuity operations are performed for debilitating and malignant conditions. A good example is pancreatic resection, in which patients' baseline physiological condition is generally suboptimal, operations are formidable, and long-term survival is rare. Until recently, elderly patients with periampullary malignant neoplasms were often precluded from undergoing resection by surgeons for these reasons, and this likely still affects referral patterns. This rationale was largely based on historical data suggesting marginal improvement in survival, coupled with operative mortality rates as high as 20%.<sup>25-27</sup> However, with global improvements in care, current recommendations for management should not consider age alone to be a contraindication for resection.<sup>28-31</sup>

Pancreatic resection, today, like other operations, can be performed safely and frequently in elderly patients.<sup>12,29,30</sup> This study demonstrates that a substantial 18% of all resections in our specialty practice were for elderly patients; however, operative mortality (1%) exceeded benchmark standards for pancreatic resection at high-volume centers (3.8%).<sup>14,32</sup> Furthermore, although the overall rate of complications was high, major morbidity was infrequent (21%). Intraoperative blood loss, rather than age, was the strongest predictor. High standards of surgical care were similarly achieved for other clinical and economic metrics: hospital stays were short, use of the intensive care unit was rare, and costs were contained.

These outcomes were scrutinized to determine whether traditional standards of quality in elderly patients rival those of younger patients. Our initial experience demonstrates that older patients have significant increases in complications, resource utilization, duration of hospital stay, and costs. However, deeper analysis—using the O:E morbidity calculations and DBCM—reveals that these outcomes depend on several factors.

The O:E analysis demonstrates that, overall, outcomes were actually slightly better than expected. Over time, surgical performance also improved considerably for both groups but was more impressive among the younger cohort. One explanation for this phenomenon is the implementation of a detailed clinical pathway and the adoption of several process and system improvements.<sup>17</sup> These initiatives, which were generally oriented to the requirements of nonelderly patients, helped reduce thromboembolic, respiratory, infectious, and intra-abdominal complications. However, the present analysis has brought to light the following 4 themes, not originally addressed in our standardized care pathway, which affect elderly patients disproportionately: postoperative delirium, blood transfusion, supplemental nutritional support, and postoperative disposition.

Analysis using DBCM demonstrates that the severity of complications also dictates clinical outcomes. Minor events, although more frequent in the elderly, had no measurable effect on clinical outcomes or costs. In this deviation category, morbidity, duration of hospital stay, and costs matched the established benchmarks of younger patients. Moderate complications resulted in more therapeutic interventions and a marginal increase in hospital stays. Major complications, however, were considerably more debilitating among older patients. Hospital stays were nearly

doubled, and elderly patients regularly required more invasive interventions to shepherd them to full recovery. This reflects early vigilance and relative aggressiveness to control the devastating effects of major complications in the elderly, who have less reserve to overcome such predicaments. That intraoperative blood loss is the only defined association with the development of major complications speaks to the importance of the original operative endeavor and suggests that specialized expertise in these operations is requisite for optimal outcomes. In sum, DBCM estimates that, overall, an elderly patient will experience a 2.2-day increase in hospital stay beyond that of younger patients, costing an additional \$9577.

There are inherent limitations to this analysis. This study was conducted within a single specialty practice at a high-volume institution, preventing comparison across the variety of practices performing these operations. Patients may differ with respect to comorbid illness, pathological disease, surgical techniques, and management strategies. Second, posthospitalization costs, particularly rehabilitation placement, likely are considerable. Therefore, the reported cost differential between elderly and nonelderly patients underestimates the overall impact. Third, we defined elderly patients as those 75 years or older. This cutoff, although seemingly arbitrary, was selected because many physicians consider patients older than 75 years to be of too high a surgical risk, particularly for invasive operations with considerable blood loss and great propensity for postoperative morbidity. Furthermore, although people 65 years or older now represent the fastest growing segment of our society, those 75 years or older will create the greatest demand for health care in the future. Little is yet known about surgical outcomes among this cohort, particularly for pancreatic cancer, for which few patients are afforded the option of surgery. Finally, people aged 65 to 75 years are increasingly regarded as surgical candidates. When successful, surgery can expect to extend these patients' lives another 10 to 15 years or more. A consolidation of people older than 65 years with those aged 65 to 75 years is appropriate because previous studies show that these age groups have similar outcomes. For these reasons, we chose an age cutoff of 75 years, although we acknowledge that a more (ie,  $\geq 80$  years) or less (ie,  $< 65$  years) stringent analysis may provide different outcomes, but likely would not significantly alter the concepts borne out herein. In addition, this study was limited to pancreatic resections because these operations represent high-acuity procedures with a significant risk of morbidity based on heterogeneous patient demographics and the natural history of the pathological disease encountered. It remains unclear whether similar encouraging outcomes will manifest in elderly patients who undergo other high-acuity operations (ie, hepatectomy, esophagectomy, and abdominoperineal resection). Nevertheless, the principles of this analysis can easily be extended to such scenarios. Finally, this retrospective analysis did not examine or compare long-term functional and quality-of-life metrics, an emerging concept deserving of detailed prospective analysis.

Despite these limitations, the findings of this study have prompted us to adopt 4 age-related process improve-

ment measures to target elderly patients at risk for hemodynamic compromise, postoperative delirium, and infectious complications. Specifically, we now use geriatric consultation preoperatively for all patients 75 years or older to address cardiorespiratory conditions, adjust plans for fluid resuscitation and pain control, and limit postoperative delirium. These dedicated gerontologists, who are familiar with complications in high-acuity surgery, provide additional evaluation of patients' cognitive and functional status. Unlike anesthesiologists and surgeons, they are trained to address important but under-recognized issues that profoundly affect surgical outcomes. These include delirium, dementia, depression, anorexia, skin breakdown, polypharmacy, glucose intolerance, and the multifactorial interactions that affect cardiorespiratory, digestive, and neurocognitive function. Other disciplines trained in the assessment and management of these issues can help achieve optimal outcomes but probably at higher cost and without access to the venues of postacute care that are more readily available to geriatricians. Therefore, we endorse that all patients undergo geriatric evaluation and comanagement, believing that this does not supplant the assessments of other specialists but instead augments optimal care.

As has been our practice, all patients, elderly and non-elderly, routinely undergo preoperative evaluation for anemia and bleeding diatheses. When abnormalities are discovered, further evaluation of causality is performed, and values are corrected before resection is performed. Transfusion is applied more liberally for elderly patients in the operative and postoperative settings. All patients 75 years or older now receive prophylactic placement of feeding jejunostomy tubes to reduce reliance on total parenteral nutrition with its attendant risks of infection and thrombosis. There have been few adverse events associated with this practice. A single elderly patient had the use of an externally damaged jejunostomy tube aborted in the early postoperative period. Other complications, such as gastrointestinal tract bleeding, obstruction, bowel ischemia, or site infections, have not been observed. Finally, elderly patients undergo early postoperative disposition planning during the initial gerontology consultation to facilitate the swifter transition from the acute-care hospital setting.

In summary, this study suggests that current standards for pancreatic resection in the elderly can and should mirror those for younger patients. However, these results should be interpreted cautiously. We treated a carefully selected population that was referred, largely by gastroenterology specialists, for surgery at a multidisciplinary specialty center, inferring an element of referral bias. Nevertheless, the results of this study should broaden current considerations for operative candidates in this domain. We believe that regular geriatric consultation, incorporated into a standardized care pathway, can be a model to improve other high-acuity elective operations for this cohort.

**Accepted for Publication:** October 15, 2008.

**Correspondence:** Charles M. Vollmer Jr, MD, Department of Surgery, Beth Israel Deaconess Medical Center, 330 Brookline Ave, Stoneman 9, Boston, MA 02215 (cvollmer@bidmc.harvard.edu).

**Author Contributions:** *Study concept and design:* Pratt, Schreiber, Lipsitz, and Vollmer. *Acquisition of data:* Pratt, Gangavati, Agarwal, Callery, and Vollmer. *Analysis and interpretation of data:* Pratt, Gangavati, Lipsitz, and Vollmer. *Drafting of the manuscript:* Pratt and Lipsitz. *Critical revision of the manuscript for important intellectual content:* Pratt, Gangavati, Agarwal, Schreiber, Lipsitz, Callery, and Vollmer. *Statistical analysis:* Pratt. *Obtained funding:* Pratt and Lipsitz. *Administrative, technical, and material support:* Agarwal, Schreiber, Lipsitz, Callery, and Vollmer. *Study supervision:* Gangavati, Lipsitz, Callery, and Vollmer.

**Financial Disclosure:** None reported.

**Funding/Support:** This study was supported by a Clinical Research Fellowship Award from the Doris Duke Charitable Foundation (Dr Pratt), the PASTEUR program (Dr Pratt) and the Office of Enrichment Programs (Dr Pratt) at Harvard Medical School, the Edyth and Irving S. Usen and Family Chair in Geriatric Medicine at Hebrew Senior Life (Dr Lipsitz), and grant PO1AG004390 from the National Institute on Aging (Dr Lipsitz).

**Additional Contributions:** Gail Piatkowski, BS, Kathleen Murray, BS, Elizabeth Wood, BS, and Kenneth Sands, MD, MPH, of the Department of Health Care Quality Outcomes at Beth Israel Deaconess Medical Center, provided support and assistance for this study.

## REFERENCES

- He W, Sengupta M, Velkoff VA, DeBarros KA; US Census Bureau. *65+ in the United States: 2005*. Washington, DC: US Government Printing Office; 2005.
- Etzioni DA, Liu JH, O'Connell JB, Maggard MA, Ko CY. Elderly patients in surgical workloads: a population-based analysis. *Am Surg*. 2003;69(11):961-965.
- National Cancer Institute. Cancer of the pancreas. Surveillance Epidemiology and End Results Web site. <http://seer.cancer.gov>. Accessed April 2, 2008.
- Michaud DS. Epidemiology of pancreatic cancer. *Minerva Chir*. 2004;59(2):99-111.
- Stojadinovic A, Brooks A, Hoos A, Jaques DP, Conlon KC, Brennan MF. An evidence-based approach to the surgical management of resectable pancreatic adenocarcinoma. *J Am Coll Surg*. 2003;196(6):954-964.
- Lowenfels AB, Maisonneuve P. Epidemiology and prevention of pancreatic cancer. *Jpn J Clin Oncol*. 2004;34(5):238-244.
- Devesa SS, Grauman DJ, Blot WJ, Pennello G, Hoover RN, Fraumeni JF Jr. *Atlas of Cancer Mortality in the United States, 1950-94*. Washington, DC: National Institutes of Health; 1999. NIH publication 99-4564.
- Neoptolemos JP, Dunn JA, Stocken DD, et al; European Study Group for Pancreatic Cancer. Adjuvant chemoradiotherapy and chemotherapy in resectable pancreatic cancer: a randomised controlled trial. *Lancet*. 2001;358(9293):1576-1585.
- Aloia TA, Lee JE, Vauthey JN, et al. Delayed recovery after pancreaticoduodenectomy: a major factor impairing the delivery of adjuvant therapy [published correction appears in *J Am Coll Surg*. 2007;204(6):1304]. *J Am Coll Surg*. 2007;204(3):347-355.
- Cameron JL, Pitt HA, Yeo CJ, Lillemoe KD, Kaufman HS, Coleman J. One hundred and forty-five consecutive pancreaticoduodenectomies without mortality. *Ann Surg*. 1993;217(5):430-438.
- Porter GA, Pisters PW, Mansyur C, et al. Cost and utilization impact of a clinical pathway for patients undergoing pancreaticoduodenectomy. *Ann Surg Oncol*. 2000;7(7):484-489.
- Balcom JH IV, Rattner DW, Warshaw AL, Chang Y, Fernandez-del Castillo C. Ten-year experience with 733 pancreatic resections: changing indications, older patients, and decreasing length of hospitalization. *Arch Surg*. 2001;136(4):391-398.
- Birkmeyer JD, Finlayson EVA, Birkmeyer CM. Volume standards for high-risk surgical procedures: potential benefits of the Leapfrog initiative. *Surgery*. 2001;130(3):415-422.
- Traverso LW, Shinchi H, Low DE. Useful benchmarks to evaluate outcomes after esophagectomy and pancreaticoduodenectomy. *Am J Surg*. 2004;187(5):604-608.
- McGory ML, Shekelle PG, Rubenstein LZ, Fink A, Ko CY. Developing quality indicators for elderly patients undergoing abdominal operations. *J Am Coll Surg*. 2005;201(6):870-883.
- Cameron JL, Riall TS, Coleman J, Belcher KA. One thousand consecutive pancreaticoduodenectomies. *Ann Surg*. 2006;244(1):10-15.
- Vollmer CM Jr, Pratt W, Vanounou T, Maithe SK, Callery MP. Quality assessment in high-acuity surgery: volume and mortality are not enough. *Arch Surg*. 2007;142(4):371-380.
- Vanounou T, Pratt W, Fischer JE, Vollmer CM Jr, Callery MP. Deviation-based cost modeling: a novel model to evaluate the clinical and economic impact of clinical pathways. *J Am Coll Surg*. 2007;204(4):570-579.
- Pratt W, Joseph S, Callery MP, Vollmer CM Jr. POSSUM accurately predicts morbidity for pancreatic resection. *Surgery*. 2008;143(1):8-19.
- American Society of Anesthesiologists. New classification of physical status. *Anesthesiology*. 1963;24(1):111.
- Karnofsky DA, Burchenal JHK. The clinical evaluation of chemotherapeutic agents in cancer. In: McLeod CM, ed. *Evaluation of Chemotherapeutic Agents*. New York, NY: Columbia University; 1948:191-205.
- Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg*. 2004;240(2):205-213.
- Copeland GP, Jones D, Walters M. POSSUM: a scoring system for surgical audit. *Br J Surg*. 1991;78(3):355-360.
- Hosmer DW, Lemeshow S. *Applied Logistic Regression*. 2nd ed. New York, NY: John Wiley & Sons Inc; 2000.
- Forrest JF, Longmire WP Jr. Carcinoma of the pancreas and periampullary region. *Ann Surg*. 1979;189(2):129-138.
- Herter FP, Cooperman AM, Ahlborn TN, Antinori C. Surgical experience with pancreatic and periampullary cancer. *Ann Surg*. 1982;195(3):274-281.
- Kairaluoma MI, Kiviniemi H, Stahlberg M. Pancreatic resection for carcinoma of the pancreas and the periampullary region in patients over 70 years of age. *Br J Surg*. 1987;74(2):116-118.
- Fong Y, Blumgart LH, Fortner JF, Brennan MF. Pancreatic or liver resection for malignancy is safe and effective for the elderly. *Ann Surg*. 1995;222(4):426-437.
- Lightner AM, Glasgow RE, Jordan TH, et al. Pancreatic resection in the elderly. *J Am Coll Surg*. 2004;198(5):697-706.
- Brozzetti S, Mazzoni G, Miccini M, et al. Surgical treatment of pancreatic head carcinoma in elderly patients. *Arch Surg*. 2006;141(2):137-142.
- Makary MA, Winter JM, Cameron JL, et al. Pancreaticoduodenectomy in the very elderly. *J Gastrointest Surg*. 2006;10(3):347-356.
- Birkmeyer JD, Siewers AE, Finlayson EV, et al. Hospital volume and surgical mortality in the United States. *N Engl J Med*. 2002;346(15):1128-1137.